

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 July 2002 (11.07.2002)

PCT

(10) International Publication Number  
**WO 02/053720 A1**

(51) International Patent Classification<sup>7</sup>: C12N 9/02,  
15/82, A01H 5/10

(21) International Application Number: PCT/IB00/02045

(22) International Filing Date:  
29 December 2000 (29.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicants: CARLSBERG RESEARCH LABORATORY [DK/DK]; Gamle Carlsberg Vej 10, DK-2500 (DK). HEINEKEN TECHNICAL SERVICES BV [NL/NL]; 2380 BB Zoeterwoude (NL). BRASSERIES KRONENBOURG [FR/FR]; 68, route d'Oberhausbergen, F-67037 Strasbourg (FR).

(72) Inventors: DOUMA, Anneke, Christiana; Eikenlaan 11B, NL-3707 SB Zeist (NL). DODERER, Albert; P.O. Box 510, NL-2380 BB Zoeterwoude (NL). CAMERON-MILLS, Varena; Kirkevaenget 20, DK-2500 Valby (DK). SKADHAUGE, Birgitte; Carlsberg Research Laboratory, Lindehoejvej 31B, DK-3460 Birkerød (DK). BECH, Lene, Molskov; Rørkaer 6, DK-2765 Smørum (DK). SCHMITT, Natalie; Cornelis

Jostraat 46, NL-2584 ET Den Haag (NL). HEISTEK, Jolanda, Carolina; Van Blankenheimstraat 15, NL-3132 VA Vlaardingen (NL). VAN MECHELEN, Johannes, Reinier; Brouwersgracht 214, NL-1013 HD Amsterdam (NL).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/053720 A1

**BEST AVAILABLE COPY**

(54) Title: LOW-LIPOXYGENASE 1 BARLEY

(57) Abstract: Barley plants having reduced lipoxigenase-1 enzyme activity are provided, for example, barley plants expressing mutant LOX-1 protein. The barley plants of the invention are useful in the production of plant products such as malt and brewed beverages, particularly beer, having increased stability and reduced T2N potential.

## LOW-LIPOXYGENASE 1 BARLEY

This application is being filed as a PCT International patent application in the name of Carlsberg Research Laboratory, a Denmark corporation, on 29 December 2000, designating all countries except the

5 U.S.

### FIELD OF THE INVENTION

This invention is in the field of plant biotechnology. More specifically, the invention relates to a mutant barley lipoxygenase 1 gene (lox-1) that encodes an enzyme with severely reduced 9-hydroperoxy-octadecanoic acid forming activity. The invention also relates to the use of barley cultivars homozygous for lox-1 in brewing processes to reduce the formation of off-flavors in brewed products, such as beer, during storage.

15

### BACKGROUND OF THE INVENTION

Lipoxygenases are a family of enzymes (EC 1.13.11.12) that catalyze the dioxygenation of free and esterified poly-unsaturated fatty acids containing a 1(Z), 4(Z)-pentadiene configuration. The products of lipoxygenase-catalyzed reactions have long been suspected as major culprits for the appearance of stale flavors in plant grain/seed and grain/seed derived food products (Robinson *et al.*, 1995, *Food Chem.*, 54: 33-43). Lipoxygenases have been implicated in the production of volatile hexanal aldehydes generated during soybean processing, which have an undesirable aroma, limiting the use of soybean proteins in food products. Three lipoxygenase isozymes expressed in soybean seed are believed to contribute to lipid oxidation and hexanal formation. Soybean mutants lacking one or more of these isozymes have been generated with the aim of reducing hexanal formation and improving their flavor stability. The success of this approach has been evaluated by Hildebrand *et al.*, 1990, *J. Agric. Food Chem.* 38: 1934-1936. Mutants lacking soybean lipoxygenase 3 produced higher hexanal levels, suggesting that

25

30

this isozyme diverts 13-hydroxyperoxyoctadecanoids, produced by lipid oxidation, towards non-volatile products. The field performance of triple-null soybean lines, lacking all three seed lipoxygenases, has shown that these enzymes are not essential for normal agronomic and seed characteristics (Narvel *et al.*, 1998, *Crop Sci.* 38: 926-928).

Lipoxygenases have also been implicated in the generation of off-flavors in rice, which can occur during grain storage. The release of free fatty acids can be detected in stored grain, which is indicative of the metabolism of the triglycerides reserves. The rice variety Daw Dam was found to accumulate lower levels of pentanals and hexanals giving a better flavor stability on storage (Susuki *et al.*, 1999, *J. Agric. Food Chem.*, 47: 1119-1124). This desirable phenotype was attributed to the absence of rice lipoxygenase-3, which oxidises unsaturated lipid acyl chains to form 9-hydroxyperoxyoctadecanoic positional isomers.

It is recognised that the lipoxygenase pathway is complex with many branches and its role in numerous aspects of plant growth and physiology are not fully understood. Modifications of the lipoxygenase pathway which alter 9-hydroperoxidation activity in seed crops are proposed to regulate their susceptibility to mycotoxin contamination by *Aspergillus* spp. (WO 9726364), which is consistent with the involvement of this pathway in plant pathogen resistance, but is not related to the aims of the invention herein described.

Among the many aroma volatiles which contribute to the flavor of beer, the higher unsaturated aldehydes with a 6-12 carbon chain have particularly low organoleptic flavor thresholds (Meilgaard 1975, *MBAA Tech. Quart.* 12: 151-168). *Trans*-2-nonenal, which is a member of this group, has both an extremely low flavor threshold of 0.11 ppb and contributes an unpleasant straw-like, "cardboard" flavor to the beer. The characteristic off-flavor caused by *trans*-2-nonenal is a common characteristic of beers stored for 1-3 months or more and is particularly detrimental to the flavor of lager beer, which is brewed with light malts and has a delicate flavor.

Sulfite has long been known to improve the flavor stability of beer, not only by binding oxygen and acting as an anti-oxidant, but also by forming volatile bisulfite addition compounds with aldehydes and ketones present in the beer. The two major sources of sulfite in beer are sulfite produced by yeast during fermentation via the sulfur assimilation pathway and sulfite added to the beer prior to packaging. Fermentation conditions that enhance yeast sulfite production and secretion will allow the formation of sulfite-carbonyl adducts from carbonyls present in the wort and prevent their further metabolism by the yeast (Dufour 1991, *Proc.Eur. Brew. Conv. Congr.*, Lisbon, pp. 209-216). In this manner carbonyls such as acetaldehyde and diacetyl may be transferred to the beer. The ability of sulfite to prevent the appearance of the carbonyl compound *trans*-2-nonenal during beer aging has been demonstrated by brewing beer with a yeast strain blocked in the sulfur assimilation pathway (Johannesen *et al.*, 1999, *Proc.Eur. Brew. Conv. Congr.*, Nice, pp. 655-662). Following bottling, the beer was subjected to forced aging by storing it at 37°C for 7 days, after which *trans*-2-nonenal levels were found to be well above the taste-threshold. If 10 ppm sulfite was added to the low-sulfite beer just prior to bottling, the appearance of *trans*-2-nonenal during forced aging was significantly reduced. The reaction between sulfite and carbonyl compounds is reversible and under thermodynamic and kinetic control. The apparent equilibrium constants for bisulfite compounds ranges from  $10^{-6}$  M for carbonyl compounds such as acetaldehyde, hexanal, and decanal, to  $10^{-3}$  for diacetyl and pyruvate (Dufour 1991, *supra*). During beer storage, gas exchange through the packaging will allow oxygen into the beer and sulfite will be lost, such that weaker bisulfite adducts will dissociate, allowing free carbonyls to appear in the beer. While sulfite unquestionably enhances the flavor-stability of beer, particularly in the short-term, its retention in packaged beer is strongly dependent on gas exchange through the packaging and temperature. In a finished beer the natural levels of sulfite produced during fermentation are variable and the addition of sulfite prior to



bottling is not a universally accepted practice. For these reasons sulfite alone does not provide a reliable method to enhance the long-term flavor-stability of beer under the different beer storage conditions used around the globe.

5           It is generally accepted that the *trans*-2-nonenal found in beer results from the oxidation of polyunsaturated fatty acids derived from barley grain lipids, where the 18-carbon chain fatty acid, linoleic acid [classified as an 18:2,n-6 polyunsaturated fatty acid (Broun, Gettner and Sommerville 1999, *Annu. Rev. Nutr.* 19: 197-216)] is the most abundant.

10          However, there is little agreement in the literature as to the mechanism whereby *trans*-2-nonenal is formed. The presence of enzymatic pathways leading to *trans*-2-nonenal formation from poly-unsaturated fatty acids has been proposed, but the individual enzymatic steps have never been demonstrated experimentally in barley grain or during the

15          malting process (Gardner 1988, *Adv. Cereal Sci. Technol.* 9: 161-215). The concept of using anti-sense or co-suppression gene technology to reduce lipoxygenase-1 levels in barley grain, and thereby control 9-hydroperoxidation and reduce aldehyde and alcohol levels in the finished barley grain, has been proposed as a means to control off-flavor

20          formation, but results of such an approach are not reported (McElroy and Jacobsen, 1995, *Bio/Technology* 13: 245-249).

          A forcing test has been developed as a method for assessing the *trans*-2-nonenal potential of a beer, where *trans*-2-nonenal formation in wort or beer is induced by subjecting samples to elevated temperatures at

25          reduced pH, (100°C, at pH 4.0 for 2 hours). Attempts to correlate the *trans*-2-nonenal potential in wort and finished beer with the total level of lipoxygenase activity in the kilned malt have indicated that lipoxygenase may contribute to the appearance of *trans*-2-nonenal in aged beer (Drost *et al.*, 1990, *J. Am. Soc. Brew. Chem.* 48: 124-131). The conclusions that

30          can be drawn from this study, however, are severely limited by the fact that the lipoxygenase activity in the barley malt was regulated at the end of the malting process by the degree of enzyme inactivation during kiln

drying. Thus, only the effect of the residual malt lipoxygenase activity on the *trans*-2-nonenal potential in the derived wort and finished beer was examined. The study failed to evaluate the lipoxygenases that catalyse the first step in the lipoxygenase enzymatic pathway in the barley grain during development and malting, and their role as determinants of *trans*-2-nonenal levels found in beer. Indeed, the absence of barley cultivars deficient in one or more lipoxygenase isoenzyme has made it impossible to provide convincing evidence for the role of the lipoxygenase pathway in barley malt in controlling the formation of *trans*-2-nonenal. Such experiments are needed to evaluate the contribution of enzymatic, as compared to auto-oxidative/chemical pathways, to the formation of *trans*-2-nonenal in beer. The brewing process involves a high temperature step of wort boiling where these non-enzymatic reactions are proposed to occur (Noël *et al.*, 1999, *J. Agric. Food Chem.* 47: 4323-4326).

## SUMMARY OF THE INVENTION

This invention provides a barley cultivar having greatly reduced lipoxygenase-1 activity. In one embodiment, the barley plants of the invention contain a mutant *lox-1* gene expressing greatly reduced levels of the isoenzyme lipoxygenase-1. In an alternative embodiment, the barley plants contain a heterologous nucleic acid sequence expressing an antisense sequence to the wild-type *lox-1*, thereby reducing the enzyme's activity.

As shown herein, malt and wort produced from the reduced lipoxygenase barley of the invention, for example, from barley cultivars homozygous for a mutant *lox-1* gene, are useful to produce beer with significantly enhanced flavor stability and reduced *trans*-2-nonenal levels, particularly under conditions known to promote the appearance of T2N. The invention demonstrates a correlation between the activity of barley malt lipoxygenase-1 to produce 9-hydroxyperoxy-octadecadienoic acids (9-HPOD), and the presence of *trans*-2-nonenal in beer. The

invention further demonstrates that the use of barley homozygous for the mutant *lox-1* gene in the brewing process improves the flavor stability of the beer, both during storage and on exposure to elevated storage temperatures. These properties enhance the quality of the beer, and are  
5 useful to extend its shelf-life and reduce the need to cool beer during transport and storage.

The invention provides barley plants and portions thereof having reduced lipoxygenase-1 activity, including barley plants expressing mutant LOX-1 protein as described herein, as well as methods for  
10 producing such barley plants, plant portions, products of the plants, and particularly malt and beer products produced from the barley plants of the invention.

#### BRIEF DESCRIPTION OF THE DRAWINGS

15 Figure 1 is a graph showing the effect of the inhibitor nordihydroguaiaretic acid (NDGA) on immuno-affinity purified lipoxygenase 1 and 2 activity from embryos of 3 day germinated barley grain.

Figure 2 is a graph showing the fresh weight of developing grain  
20 of Line G and cv Vintage from 5 days after flowering to full-maturity (FM). Each determination is the mean single grain weight from 6 spikes.

Figure 3 is a graph showing the dry weight of developing grain of Line G and cv Vintage from 5 days after flowering to full-maturity (FM). Each determination is the mean single grain weight from 3 samples of 5  
25 grain.

Figure 4 is a graph showing total lipoxygenase activity in developing grain of Line G and cv Vintage from 5 days after flowering to full-maturity (FM).

Figure 5 is a graph showing 9- and 13-HPOD products of linoleic  
30 acid oxidation by lipoxygenase activity in developing grain of Line G.

Figure 6 is a graph showing total lipoxygenase activity in embryos of germinating grain of Line G and cv Vintage expressed as  $\mu$  mol/min/10 embryos (U/10 embryos).

Figure 7 is a graph showing 9- HPOD and 13-HPOD products of linoleic acid oxidation by lipoxygenase activity in embryos of germinating grain of Line G and cv Vintage, showing levels of 9- HPOD and 13-HPOD.

Figure 8 is a Western blot showing immunodetection of lipoxygenase 1 in embryos of developing grain of Line G and cv Vintage [wt] from 5 days after flowering to full-maturity (FM).

Figure 9 is a Western blot showing immunodetection of lipoxygenase 1 in embryos of grain of Line G and cv Vintage [wild-type] germinated for 0 - 6 days.

Figure 10 is a Northern blot probed with the 3' non-transcribed region of the *lox-1* cDNA and showing lipoxygenase 1 transcripts detected in developing grain of Line G and cv Vintage [wild-type] from 5 days after flowering to full-maturity (FM).

Figure 11 is a Northern blot probed with the 3' non-transcribed region of the *lox-1* cDNA and showing lipoxygenase 1 transcripts detected in embryos of grain of Line G and cv Vintage [wt] germinated for 0 - 6 days.

Figures 12A-12G are a nucleotide sequence alignment of the promoter and transcribed region of the *lox-1* wild-type cv Vintage allele (WT) and the Line G allele (LG). The transcription start site (+1), ATG start codon (+69) and translation stop codon (+4231) in the gene sequences are underlined. Nucleotide mutations identified in the Line G allele are shown in bold italics and indicated by an asterisk.

Figure 13 is a schematic presentation of the *lox-1* gene of cv Vintage (wild-type) and the mutant *lox-1* gene of Line G. The transcript from +1 to +4375 is composed of 7 exons (stippled boxes) and 6 introns (white boxes). Two mutations in the *lox-1* gene are indicated.

Figure 14 is a schematic drawing of gene cassettes for transient expression of the wild-type *lox-1* cDNA and *lox-1* gene and the mutant *lox-1* gene from Line G. The lipoxygenase coding sequences were cloned between the constitutive maize ubiquitin promoter with intron 1 (*Ubi-1*) and the *nos* terminator.

Figure 15 is a bar graph showing Lipoxygenase 1 activity in barley aleurone protoplasts transfected with gene cassettes containing the wild-type *lox-1* cDNA; the mutant *lox-1* gene from Line G; WT *lox-1* gene; and a control GUS reporter gene. Lipoxygenase activity in extracts of transfected protoplasts was assayed in microtiter plates by the oxidation of KI and quantitated spectrophotometrically. Lipoxygenase 1 activity was expressed as units per  $\mu\text{g}$  protein in the extract and is shown as the mean of 3 measurements from 2 replicate assays.

Figure 16 is a sequence alignment demonstrating that a RFLP between the wild-type and mutant *lox-1* gene is due to a point mutation at nucleotide 2347, creating an additional *AatII* restriction site.

Figure 17 is a schematic presentation of the *lox-1* PCR fragments amplified and cleaved in the polymerase chain reaction – cleavage amplified polymorphic site (PCR-CAPS) assay. The positions of PCR primers are indicated by arrows and the *AatII* sites are shown above the gene (sequence position). The exon and intron regions within the PCR product are distinguished by stippled and white boxes respectively, and the sizes of the *AatII* digestion fragments are given.

Figure 18 is an electrophoretic agarose gel showing *lox-1* PCR fragments (652 bp) amplified in the first step of the PCR-CAPS assay from Line G and cv Vintage genomic DNA.

Figure 19 is an electrophoretic agarose gel showing RFLP detected by PCR-CAPS in the wild-type and mutant *lox-1* gene. The *AatII* digestion fragments of the mutant gene include a unique 313 bp restriction fragment, indicated by an asterisk.

Figure 20 is a table showing a back-crossing program for the single recessive gene pair *ll* (low lipoxygenase trait) of Line G to cv

Alexis. The *LL* genotype are plants expressing wild-type lipoxygenase activity (dominant allele), the *ll* genotype are plants expressing the low-lipoxygenase (recessive allele). *Ll* are heterozygous plants containing both the wild-type and the low-lipoxygenase allele. Since the low-lipoxygenase trait is a recessive trait, *Ll* plants show wild-type lipoxygenase activity. After each round of back-crossing (including self-pollination), the *ll* progeny is expected to represent 25% of the progeny. The observed frequencies of low-lipoxygenase activity are indicated. The calculated percentage of the cv Alexis genetic background having the homozygous low-lipoxygenase allele is indicated as % Alexis.

Figure 21 is an electrophoretic agarose gel showing PCR-CAPS detection of the mutant *lox-1* gene in *ll* progeny of the Line G – Alexis back-cross program. PCR-CAPS assay on genomic DNA of Line G (Lane 2), cv Vintage (Lane 3), *ll* progeny of 3<sup>rd</sup> (Lane 4) and 4<sup>th</sup> back-cross (Lanes 5 – 9). DNA ladder (Lane 1). Control, backcrossed high lox line (lane 10).

Figures 22A-22B are a comparative alignment of amino acid sequences of soybean lipoxygenases LOX-1 (Gm1), LOX-2 (Gm2), LOX-3 (Gm3), and barley lipoxygenases LOX-1 (Hv1) and LOX-2 (Hv2).

## DETAILED DESCRIPTION OF THE INVENTION

In accordance with the subject invention, plant materials, plant products, and methods are provided for producing a beverage, such as beer, the beverage having a reduced content of the off-flavor compound *trans*-2-nonenal, such that the flavor stability of the beverage, e.g., beer, during storage and on exposure to elevated temperatures is improved, relative to a control beverage. More particularly, the invention provides barley varieties whose developing and germinating grain produce greatly reduced activity levels of the enzyme lipoxygenase-1, denoted LOX-1, which, for example, when used in a beer brewing process, results in a

beer having reduced *trans*-2-nonenal levels, as compared with a control barley variety.

The methods used to generate, characterize, and validate a barley variety having greatly reduced LOX-1 activity, and use of this type of  
5 barley for the production of flavor-stable beer are described below.

## 1. Definitions

As used herein, the following terms have the indicated definitions:

10 "Plant portion" means a plant or specific part of a plant, such as the stem, leaves, roots, flowers, seeds, grains, fruits, or buds.

"LOX-1" means lipoxygenase-1 protein; "*lox-1*" means the gene encoding LOX-1.

"Mutant barley *lox-1*" means a mutagenized barley gene encoding  
15 a mutant lipoxygenase 1 polypeptide.

"Non-mutated control" means a plant, nucleic acid, gene, polypeptide, plant portion, or plant product containing wild type gene or protein.

"Heterologous" means a non-native sequence, e.g., a sequence  
20 derived from another species, or a recombinantly engineered or synthetic sequence that differs from the native sequence.

"Plant product" means a product resulting from the processing of a plant or plant portion, and includes, for example, malt and wort.

"Acidic amino acid" means aspartic or glutamic acid.

25 "Basic amino acid" means histidine, lysine, or arginine.

"Polar amino acid" means threonine, serine, tyrosine, tryptophan, asparagine, or glutamine.

"Organoleptic properties" means properties appealing to the olfactory and taste senses that are analysed, for example, by a trained  
30 taste panel.

"Brewed product" means a product prepared by mashing, boiling, and fermenting, e.g., beer.

"Reduced *trans*-2-nonenal" means less than about 50%, as compared with wild-type (control) conditions.

## 2. Lipoxygenase Activity

5 Lipoxygenase enzymes catalyze the oxidation of polyunsaturated fatty acids. In barley, the isoenzymes LOX-1 and LOX-2 are known. LOX-1 primarily catalyzes 9-hydroperoxidation, whereas LOX-2 primarily catalyzes 13-hydroperoxidation of polyunsaturated octadecanoic fatty acids. The data shown in the Examples below  
10 demonstrates a correlation between barley LOX-1 9-hydroperoxidation activity and the presence of *trans*-2-nonenal in beer. Accordingly, barley having reduced LOX-1 activity is useful to produce beer having a reduced *trans*-2-nonenal level and/or potential as compared with a control.

15

## 3. Production of low lipoxygenase barley

A variety of known genetic approaches can be used to produce the plants of the invention, that is, to reduce the level of lipoxygenase 1 enzyme activity expressed in a barley plant in a stable, inheritable  
20 manner. These approaches include, but are not restricted to antisense technology and mutagenesis, such as chemical and radiation induced mutagenesis, as well as site-directed mutagenesis.

**Barley transformation.** Barley can be transformed with various  
25 nucleic acid molecules designed to manipulate *lox*-1 gene expression or alter the architecture of the *lox*-1 gene. Various methods, for example, *Agrobacterium tumefaciens*-mediated transfer (Tingay *et al*, 1997, *Plant J.*, 11: 1369-1376), particle bombardment (Wan and Lemaux, 1994, *Plant Physiol.*, 104: 37-48, or polyethylene glycol (PEG)-mediated DNA  
30 uptake (Funatsuki and Kihara, 1995, *Theor. Appl. Genet.*, 91:707-712), can be used to successfully introduce nucleic acids into a barley cell, for example into a protoplast, callus, or an embryo.



Various promoters can be used to drive expression of the gene of interest. For expression of *lox-1*-containing vectors, including antisense sequences, the native *lox-1* promoter region can be used. The promoter sequence of *lox-1* is contained in nucleotides 2602 – 3511, which  
5 includes the 5' UTR of EMBL accession no. U83904. Alternatively, promoters that drive expression of the gene of interest constitutively, for example the *Ubi.1* maize ubiquitin promoter, can be used (Wan and Lemaux, *Supra*; Kjærulff et al., in P. Mathis, Ed., 1995, *Photosynthesis: from Light to Biosphere*, Vol. II, 151-154). Expression vectors can also  
10 contain a transcription termination region, for example, the 3' terminator of the nopaline synthase gene (3'-*nos*) (Bevan, *et al*, 1983, *Nucl. Acids Res.*, 11: 369-385) has been fused to genes expressed in transgenic barley (Wan and Lemaux, *Supra*; Funatsuki and Kihara, *Supra*).

Expression vectors can also contain a gene that allows for  
15 selection of transformed cells when the vector has been successfully integrated in the cell. These genes can encode antibiotic or herbicide resistance genes, for example the neomycin phosphotransferase (*npt*) or the phosphinothricin acetyl transferase (*bar*) gene. When expressed, such resistance genes allow for growth of the transformed cell in neomycin -  
20 or bialaphos-containing media, respectively (See, for example, Wan and Lemaux, *Supra*; Funatsuki and Kihara, *Supra*; Kjærulff et al., in P. Mathis, *Supra*).

Following transformation, cells can be grown in selective media for a period of time and then cultured to allow for the formation of  
25 shoots, followed by root systems, and then plantlets. A successful barley transformation procedure was developed by Funatsuki and Kihara, (*Supra*), where transformation of barley protoplasts by PEG with neomycin phosphotransferase-containing expression vectors and subsequent selection in neomycin yielded fertile plants containing the  
30 transgene. The transgene was shown to integrate into the genome and most of the transgenic plants expressed the protein encoded by the

transgene. These transgenic plants also were able to transmit and express the transgene following crosses.

It is understood that a variety of transformation methods, expression vectors, promoters, selectable markers, and the like are known and useful for transformation of barley.

**Barley mutagenesis.** The *lox-1* gene can be targeted for site-specific mutagenesis using chimeric RNA/DNA oligonucleotides. These chimeric RNA/DNA oligonucleotides have been shown to successfully introduce mutations in plant cells (Zhu et al., 1999, *Proc. Natl. Acad. Sci.* 96: 8768-8773; and Beetham et al., 1999, *Proc. Natl. Acad. Sci.* 96: 8774-8778) and mammalian cells (Yoon et al., 1999, *Proc. Natl. Acad. Sci.* 93:2071-2076) at desired locations. The chimeric RNA/DNA oligonucleotides can be transformed into the barley protoplasts or cells of interest in a variety of ways, for example using the PEG-mediated or particle bombardment-mediated transformation methods described above. The individual protoplasts or cells can then regenerated by tissue culture to whole fertile plants, and the mutational event can be confirmed and followed, for example using a PCR-based approach as detailed in the Examples below.

This site-directed mutagenesis method can be applied to mutate specific residues in the *lox-1* gene. The *lox-1* gene can be mutated at one or more nucleotide position in the promoter region to downregulate or abolish *lox-1* transcription. Specific mutagenesis can also be applied to introduce changes in the *lox-1* coding region that, for example, reduce the enzyme's activity. Such mutations include, but are not limited to, insertions, deletions, and substitutions resulting in a frameshift, truncation of the LOX-1 protein, and/or alteration of the neutral and hydrophobic nature of the enzyme's substrate cavity.

**Antisense expression.** Reduction in *lox-1* expression can also be accomplished by expression of a *lox-1* antisense construct in the barley cells. Methods for the expression of antisense constructs in barley to reduce the expression of a targeted protein have been reported, for

example, in Gilpin, M.J. et al., 1998, In: *Photosynthesis: Mechanisms and Effects*, G. Garab, ed., Vol. IV, 2983-2986; Kjærulff et al., 1995, In: *Photosynthesis: from Light to Biosphere*, P. Mathis, Ed., Vol. II, 151-154.

5           Barley cells can be transformed with an expression construct containing an antisense nucleic acid sequence. The expression construct produces an antisense RNA molecule capable of specifically binding to at least a portion of the mRNA produced from the wild type *lox-1* gene, through complimentary base pairing, and capable of disrupting the  
10       splicing of the pre-mRNA or translation of this mRNA. A constitutive or tissue/temporal specific promoter, for example, the barley *lox-1* promoter described above, can drive expression of the antisense nucleic acid sequence.

**Chemical mutagenesis.** The chemical mutagen sodium azide  
15       ( $\text{NaN}_3$ ) has commonly been used for barley mutagenesis and is known to induce stable mutations in the DNA (deoxyribonucleic acid) sequence of the barley genome (Olsen *et al.*, 1993, *Proc. Natl. Acad. Sci. USA*, 90: 8043-8047). Other chemical mutagens, for example, ethyl  
          methanesulfonate (EMS), azidoglycerol (AG, 3-azido-1,2-propanediol),  
20       methyl nitrosourea (MNU), and maleic hydrazide (MH) can also be used to induce DNA mutations (Rank, J. et al., 1997, *Mutat. Res.* 390:121-7), as can UV irradiation.

          As shown in the Examples below, the grain of the barley cultivars (cv) Vintage and Caruso were treated with sodium azide and propagated  
25       by self-fertilization through to the 3<sup>rd</sup> generation (M3).

#### 4. Identification and Selection of Low Lipxygenase Barley

Identification and selection of barley plants having reduced lipxygenase isoenzyme activity in the grain can be achieved, for example, by analysis of lipxygenase activity. Enzymatic assays can be used to determine the activity of the two major lipxygenases known to be present in either mature or germinating grain, LOX-1 and LOX-2. Such assays should distinguish LOX-1 activity from that of LOX-2.

One selective assay of LOX-1 and LOX-2 is based on the oxidation of a poly-unsaturated fatty acid by lipxygenase and the spectrophotometric detection of the hydroperoxide product of such oxidation. The specificity of this assay for LOX-1 takes advantage of the comparative insensitivity of LOX-1 to an inhibitor, for example, NDGA, relative to LOX-2.

Selective assay can also be achieved using immunoprecipitation to selectively remove LOX-1 or LOX-2 from the assay. Specific anti-LOX-1 and anti-LOX-2 antibodies, for example, monoclonal antibodies, can be prepared from purified LOX-1 or LOX-2 as described in Holtman et. al, 1996, *Supra*.

These assay methods can be adapted for microtiter plate assay procedures, or other known repetitive, high throughput assay formats, allowing the rapid screening of many samples. These assays can be validated for screening leaf tips of germinating grain in a non-destructive manner, such that seedlings selected in the screen can be further propagated.

The loss of LOX-1 activity in putative mutants can be confirmed by assay of enzymatic activity. For example, grain extracts can be incubated with linoleic acid and the oxidation products of linoleic acid analyzed, for example, by reverse phase HPLC. The relative amounts of 9-HPOD and 13-HPOD formed from linoleic acid provides a measure of LOX-1 activity, whose major product is 9-HPOD.

As shown in the Examples below, approximately 20,000 grain of the M3 generation of mutagenized cv Vintage and cv Caruso were screened for LOX-1 and LOX-2 activity by oxidation assay in the presence of inhibitor and also by immunoprecipitation assays. Using  
5 these screening methods, a mutant in cv Vintage was found having a major reduction in LOX-1 activity, and was denoted Line G. The mutant phenotype was inherited in the M4 and M5 generations.

Seed produced from the Line G barley was deposited on December 19, 2000 with the National Collections of Industrial, Food and  
10 Marine Bacteria (NCIMB), 23 St. Machar Drive, Aberdeen, AB243RY, Scotland, UK, under the terms of the Budapest Treaty, as Accession Number: \_\_\_\_\_.

## 5. Genetic Sequences

15 A precise description of the genotypic alteration that accounts for the low-lipoxygenase phenotype in barley plants of the invention is useful for identifying plants having this genetic alteration and for crossing this genetic character into other barley cultivars in a breeding program. A variety of known molecular and biochemical methods can be  
20 used to determine the genetic basis for the low lipoxygenase phenotype.

It is generally recognized that both cis-acting and trans-acting genetic sequences can determine the expression of a given gene in the genome and the activity of the gene product. Control points in gene expression include the regulation of the timing, tissue-specificity and rate  
25 of gene transcription, the stability of the transcript and the rate of transcript translation. Both the level of gene expression and the stability and specific activity of the encoded enzyme will determine the level of enzyme activity detected in a tissue.

Alterations in a plant gene sequence can be determined by DNA  
30 sequencing of known relevant parts of the genome, while Northern analysis provides a tool to monitor stable transcript levels in a given plant

tissue. Enzyme expressed in plant tissue can be evaluated by extracting the enzyme from the tissue and measuring the enzymatic activity.

As shown in the Examples below, the identity of the genetic changes that determine the low-lipoxygenase phenotype of the Line G mutant induced in cv Vintage were determined in the following manner. The structural gene encoding the LOX-1 protein, both in the parent cv Vintage and in the Line G, was amplified by the polymerase chain reaction (PCR), and the upstream promoter sequences, which regulate expression of the gene, as well as the entire coding sequence, comprising intron and exon sequences, were sequenced.

Comparison of the nucleotide sequences of the *lox-1* gene from Line G and from wild-type cv Vintage revealed 2 nucleotide substitutions in 2 exons, of which one (at position + 2347) led to a non-conservative amino acid substitution (Glycine<sup>368</sup> → Aspartate) in the expressed protein.

Figure 22 shows an alignment of soybean (Gm: *Glycine max* L) lipoxygenases LOX1 (Acc. No. P08170), LOX2 (Acc. No. P08170), LOX3 (Acc. No. AAB41272) and barley (hv: *Hordeum vulgare*) lipoxygenases LOX1 (Acc.No. P29114) and LOX2 (Acc. No. AAB70865.1). Conserved amino acid residues and conservative substitutions of charged residues are shown in bold. Secondary structure assignments for LOX3 of soybean *Glycine max*, where H=alpha helices and E=Beta strands, are shown above the alignment, and residues relevant to enzyme function (identified by an asterisk or filled circle) are shown, as described in Skrzypczak-Jankun *et.al.*, 1997, *Proteins* 29:15-31.

Amino acid residues that participate in non-heme iron binding or essential for catalysis (\*) in soybean LOX3 include: H<sub>518</sub>, H<sub>523</sub>, H<sub>709</sub> [3 N atoms]; N<sub>713</sub>, I<sub>837</sub>. The equivalent residues in barley LOX 1 are H<sub>517</sub>, H<sub>522</sub>, H<sub>708</sub>, N<sub>712</sub>, and I<sub>862</sub>. Residues in soybean LOX3 with a predicted role in catalysis(•) are: H<sub>266</sub>, H<sub>513</sub>, H<sub>776</sub>, F<sub>264</sub>, F<sub>272</sub>, F<sub>714</sub>, W<sub>519</sub>, R<sub>552</sub>, R<sub>726</sub>, D<sub>766</sub>, D<sub>779</sub>, K<sub>278</sub>. The equivalent residues in barley LOX1 are: H<sub>261</sub>, H<sub>512</sub>, H<sub>775</sub>, F<sub>259</sub>, F<sub>267</sub>, F<sub>713</sub>, W<sub>518</sub>, R<sub>551</sub>, R<sub>725</sub>, D<sub>778</sub>, and K<sub>273</sub>.

Proline (P<sub>86, 109, 167, 171, 223, 234, 291, 311, 324, 343, 345, 371, 381, 382, 486, 541, 548, 600, 616, 627, 685, 726, 734, 788, 829, 833, 839, 857</sub>) and glycine (G<sub>49, 67, 68, 70, 91, 107, 137, 187, 192, 210, 217, 218, 260, 306, 307, 336, 392, 409, 458, 474, 490, 569, 607, 674, 676, 720, 736, 783, 828, 850, 855</sub>) residues (+) located in loops and helix-capping positions in protein secondary structures, may facilitate sharp turns and folding of the peptide backbone.

Alignment of related plant lipoxygenases indicated that the Glycine-368 in barley LOX-1 is strongly conserved. Furthermore, this residue, which corresponds to Glycine-353 in soybean LOX-1, is one of 35 highly-conserved residues out a total of 58 residues that line the substrate cavity II of the enzyme, as seen from its crystal structure. These conserved residues are highlighted (boxes) in the alignment of plant lipoxygenase sequences shown in Figure 22 (Minor *et.al.*, 1996, *Biochemistry* 35:10687-10701), and include the following barley LOX-1 residues: Y<sub>224</sub>, L<sub>268</sub>, W<sub>355</sub>, E<sub>364</sub>, G<sub>368</sub>, V<sub>369</sub>, N<sub>370</sub>, I<sub>374</sub>, L<sub>424</sub>, L<sub>499</sub>, K<sub>501</sub>, A<sub>502</sub>, V<sub>504</sub>, D<sub>508</sub>, S<sub>509</sub>, H<sub>512</sub>, Q<sub>513</sub>, L<sub>514</sub>, H<sub>517</sub>, W<sub>518</sub>, H<sub>522</sub>, I<sub>556</sub>, L<sub>559</sub>, A<sub>560</sub>, L<sub>564</sub>, I<sub>565</sub>, I<sub>570</sub>, T<sub>574</sub>, S<sub>585</sub>, Q<sub>715</sub>, Y<sub>718</sub>, N<sub>724</sub>, R<sub>725</sub>, P<sub>726</sub>, T<sub>727</sub>, L<sub>772</sub>, and I<sub>862</sub>. All but 7 of the 35 conserved residues are neutral or hydrophobic residues. The substitution of a charged residue at position Glycine – 368 in barley or at another conserved neutral or hydrophobic residue lining the substrate cavity II, is likely to disturb the structural and functional properties of the enzyme. The G→D<sub>368</sub> mutation in barley Line G LOX1(♦) is located between alpha-helix H6 and beta-strand E12.

As shown in Figure 22, the lipoxygenase family of enzymes shares a high degree of sequence conservation, which is reflected in their conserved secondary structure, determined for several members of the plant lipoxygenase family including soybean LOX1 and LOX3 (Skrzypczak-Jankun *et al.*, 1997, *supra*). Barley LOX1 shares 56.9% sequence identity and 67.8% sequence similarity with soybean LOX3. Several amino acid residues in the soybean LOX3 isoenzyme have been identified as ligands for the non-heme iron, or are suggested to be essential for its activity (denoted by \* ♦). In view of the high sequence

conservation between the barley LOX1 and the soybean LOX3, it is reasonable to predict that residues in the barley LOX1 sequence that are homologous to those identified as important for the function of LOX3 may also be essential for enzymatic activity. Thus, non-conservative amino acid substitutions at any of these positions, including substitutions of those residues in barley LOX1 corresponding to the 35 highly conserved residues of soybean LOX3 that line the substrate cavity and in other positions essential for enzyme activity, are likely to reduce lipoxygenase activity.

10           The amino acid residues proline and glycine are known to facilitate turns in a peptide backbone when they are located between secondary structural elements, which allow a protein to assume a folded tertiary structure. Proline and glycine residues are also common in helix capping motifs (Parker and Hefford, 1997, *Protein Eng.*, 10: 487-496, <http://www.expasy.ch>). The single non-conservative substitution in Line G LOX1, where a glycine located between two predicted structural elements was replaced by aspartate, led to a significant loss of enzyme activity. It is thus predicted that mutation in the *LOX-1* gene causing a non-conservative amino acid substitutions at one or more of the proline or glycine residues in the barley LOX1, located in regions outside the structural elements, may similarly prevent folding of the native protein and consequently reduce the activity of the encoded enzyme.

          Thus, in one embodiment, a useful mutant barley plant of the invention having reduced lipoxygenase 1 activity contains a mutated nucleic acid sequence that alters the neutral or hydrophobic nature of the substrate cavity of the enzyme by insertion of one or more acidic, basic, or polar amino acids. For example, a useful nucleic acid sequence [SEQ ID NO: 11 encodes a barley LOX-1 protein [SEQ ID NO: 12] having a substitution at amino acid 368 from Glycine to Xaa, where Xaa is an acidic, basic, or polar amino acid. One specific amino acid sequence of the barley mutant LOX-1 of the invention is that where Xaa is aspartic acid, e.g., Line G.



As shown in the Examples below, the genotypic changes in Line G had no detectable influence on *lox-1* gene expression, but the LOX-1 activity detected in mature and germinating grain of Line G were approximately 9% of that detected in grain of the parent line, cv Vintage.

5 In order to provide direct evidence that the amino acid mutation in LOX-1 of Line G was responsible for the low-LOX-1 phenotype, the coding sequence of Line G *lox-1* and cv Vintage *lox-1* were expressed transiently in protoplasts from barley aleurone, and the activity of the mutant LOX-1 enzyme was shown to be strongly reduced in comparison to the wild-type  
10 LOX-1 enzyme.

#### 6. Transfer between breeding lines

The detection of alterations in genetic character of the barley plants of the invention genotype is useful to identify the presence of a  
15 specific genetic character in a barley line, and to facilitate the transfer of this character between breeding lines in a breeding program. A variety of molecular tools are available for the detection of alterations in genomic sequence. Such methods include, but are not restricted to, detection of restriction fragment length polymorphisms (Gebhardt and  
20 Salamini 1992, *Int. Rev. Cytology.*, 135: 201-237) and quantitative PCR based detection methods such as amplification using fluorescent primers, e.g. the TaqMan primer probe systems (Ibrahim *et al.*, 1998, *Anal. Chem* 70, 2013-2017). The choice of detection method will depend on the specific genetic character but should preferably be rapid and provide  
25 clearly interpretable data.

As shown in the Examples below, a PCR-Cleavage Amplified Polymorphic Site assay (PCR-CAPS) was provided for the detection of the mutant lipoxygenase-1 gene of Line G. The nucleotide substitution in the *lox-1* gene in Line G at position + 2347 introduced an additional  
30 site of recognition by the *AatII* restriction endonuclease that can be detected by the PCR-CAPS assay. Suitable detection methods for *lox-1*

are not restricted to this assay, but can equally well be based on TaqMan technology, and other known detection methods.

Also shown in the Examples below, the PCR-CAPS assay was applied to 4 generations of breeding material from a back-cross program, where the low-lipoxygenase phenotype in Line G was systematically back-crossed into cv Alexis. Inheritance of the low-lipoxygenase phenotype was shown to follow the inheritance of the *lox-1* gene, and the phenotype was identified as recessive and only seen in lines homozygous for the *lox-1* gene.

Accordingly, plant progeny of the invention includes breeding lines, for example, derived in a back-crossing program, that contain mutant *lox-1* and express a low lipoxygenase phenotype.

## 7. Brewing

The barley plants of the invention, including plant parts, plant progeny, grain, and plant products such as malt and wort, having low lipoxygenase 1 activity, are demonstrated herein to be useful for the manufacture of a beverage having reduced levels of free *trans*-2-nonenal over a measured period of time, or under conditions of elevated storage temperature, as compared to a beverage produced from a wild-type control barley variety. For the purpose of these comparisons the sulfite content of the beer is controlled to 5ppm or below, since it is recognised that higher sulfite levels at the time of bottling will temporarily delay the appearance of free *trans*-2-nonenal. For example, beer brewed from malt derived from the mutated barley Line G described herein, possessed stabilized organoleptic properties over a measured period of time as compared with beer brewed from malt derived from a control, non-mutated barley.

Brewing trials and evaluation of bottled beer provide the best method for evaluating the influence of different ingredients on the quality and stability of the finished beer. In order to test the influence of different barley malts, sufficient barley grain is needed to perform the

malting and brewing trials on a pilot scale and semi-industrial scale. During the period of barley propagation, the field performance of the barley line can be evaluated. The malting properties of a barley line can be evaluated during pilot or industrial scale malting, and should

5 preferably lie within national malting quality recommendations eg. the European Brewing Convention recommendations for malting quality (Analytica-EBC/European Brewing Convention, 1998, Publ. Hans Carl Getränke-Fachverlag, Nürnberg, Germany). Following pilot or semi-industrial scale brewing, the beer is packaged in brown bottles and cooled

10 to 5°C for optimal storage. At this stage the fresh beer can be analysed by trained taste panels able to detect specific beer flavors, including the off-flavor compound *trans*-2-nonenal. Additionally, the beer is chemically analysed for major flavor components including *trans*-2-nonenal. These methods of beer quality analysis are then repeated on the

15 beer following various storage conditions known to reveal the long-term storage stability of the beer, for example, forced aging treatments.

As shown in the Examples below, Line G barley was propagated in the field over several seasons in order to malt 10 tons of this line in an industrial malthouse. The control barley varieties cv Vintage and cv

20 Nevada, both having the wild-type LOX-1 phenotype, were malted under similar conditions. The kilned malt from Line G and the control barley cultivars lay within the specifications required for the semi-industrial brewing trials.

Brewing trials were performed on a 30-hl scale and evaluation of

25 the freshly bottled beers revealed that beers brewed from malt of both Line G and the control cultivars had a *trans*-2-nonenal content below the taste-threshold and were deemed satisfactory by a taste-panel. Two forced-aging treatments, either storage at 37°C for 7 days or storage for 6 to 12 weeks at 30°C, were used to evaluate the flavor-stability of the beer.

30 The flavor-stability of beer brewed from Line G malt were found to be superior to that of control malt, both with respect to taste panel

evaluation as well as the level of free *trans*-2-nonenal, and the improvement was found to be statistically significant.

## EXAMPLES

5           The present invention is further defined in the Examples below. It should be understood that the Examples, while indicating preferred embodiments, are given by way of illustration only.

### Example 1

#### 10           **Screening and Selection of Lipoxygenase Isoenzyme** **Mutants from Mutagenised Barley**

##### **1. Barley mutagenesis**

Grains of barley, *Hordeum vulgare* cv Vintage and cv Caruso,  
15       were mutagenised with sodium azide according to a published procedure (Kleinhofs *et al.*, 1978 *Mutation Research* 51: 29-35). The mutagenesis introduces point mutations in the genomic DNA that, for example, may result in amino acid changes in encoded proteins. The mutated M1 grains were propagated in the greenhouse through two generations, and  
20       the M3 grain collected for screening. The observed frequency of single gene trait mutants in the M2 generation, according to Kleinhofs *et al.*, 1978, *supra*, are 1.0-2.7 mutants per 10,000 grain from the M2 generation. Since most gene mutations are recessive and only detectable in the homozygous state, the mutagenized population was screened at the  
25       M3 generation where the expected proportion of homozygous mutant grain would be higher. A mutation frequency of 0.9 – 2.3 per 10,000 grain was expected in the mutagenized material at M3.

##### **2. A non-destructive assay of lipoxygenase 1 (LOX-1) and** 30       **lipoxygenase-2 (LOX-2) activity in M3 mutagenized grain**

A rapid screening procedure for detection of mutant barley grain with reduced LOX-1 activity was developed with the following criteria:

The screening procedure should not prevent propagation of the grain/seedling; the selected grain/seedling tissue should express quantifiable levels of lipoxygenase activity; the assay should distinguish LOX-1 activity from that of LOX-2; and the assay procedure should encompass multiple samples.

The levels of total lipoxygenase activity in different tissues of the germinating grain, namely the shoot, root, and scutellum tissue of embryo and the endosperm were assayed as follows: Extracts of barley seedling tissue were prepared by homogenising the tissue in ice-cold 20 mM Tris-HCl, pH 7.5, containing 2 mM NaN<sub>3</sub> and 0.5 mM phenylmethylsulfonyl fluoride (PMSF), followed by removal of insoluble material by centrifugation at 1000 g for 10 minutes. Lipoxygenase activity in 100 µl extract was assayed at 25°C, by addition of 2.9 ml of 20 mM linoleic acid substrate, prepared by dispersing 35 µl linoleic acid (free acid, L-1376, Sigma, USA) in 5 ml H<sub>2</sub>O containing 1% Tween 20. The reaction was followed spectrophotometrically, where the rate of increase in absorbance at 234 nm ( $A_{234\text{nm}}$ ), due to the formation of conjugated diene in the hydroperoxide product, is proportional to the enzyme activity present. One unit of lipoxygenase activity is defined as  $\Delta A_{234} = 0.001$  per minute in a 3-ml reaction, equivalent to the oxidation of 0.12 µmole linoleic acid.

The leaf tissue of grain germinated for 4 days in the dark had the highest detected levels of lipoxygenase activity (Holtman *et al.*, 1996, *Plant Physiology* 111: 569-576). Leaf tips from 4-day seedlings were thus selected for the non-destructive lipoxygenase screening assay. The pH optimum of total barley lipoxygenase activity was tested between pH 4.5 and pH 9.0 and found to be pH 6.5. Hence a 25 mM HEPES buffer (pH 6.5) containing 0.2 M boric acid was selected for the screening assay.

Since both LOX-1 and LOX-2 enzymes were immuno-detected in shoots of 4-day seedlings (Holtman *et al.*, 1996, *supra*), a LOX-1 and LOX-2 specific assay was used. The lipoxygenase inhibitor

nordihydroguaiaretic acid (NDGA), identified by Eskin *et al.*, 1977, *Crit. Rev. Food, Science and Nutrition* 9: 1-40, was found to be a selective inhibitor of barley lipoxygenases. NDGA at  $1 \times 10^{-5}$  M strongly inhibited purified barley LOX-2, while LOX-1 retained 47 % activity (Figure 1).

- 5 The selectivity of this inhibitor was tested in the leaf tip assay, by determining the ratio of 9-hydroperoxyoctadecanoid (9-HPOD) to 13-hydroperoxyoctadecanoid (13-HPOD), which result from linoleic acid oxidation by LOX-1 and LOX-2, respectively. In the lipoxygenase assay of cv Vintage leaf tips, the proportion of 13-HPOD formed fell from
- 10 24.5% to 9.5% on addition of  $1 \cdot 10^{-5}$  M NDGA.

- A selective assay for LOX-2 activity in leaf tip extracts was based on the use of LOX-1-specific monoclonal antibody (5D2) (Holtman *et al.*, 1996, *supra*) to immunoprecipitate LOX-1 present in the extracts. The residual lipoxygenase activity detected in the extracts after LOX-1
- 15 precipitation provided a measure of LOX-2 activity. The efficiency of this immunoprecipitation (described below) was evaluated by quantifying the residual LOX-1 and LOX-2 in the extract supernatant by ELISA assay, using specific monoclonal antibodies against LOX-1 (denoted 5D2) and LOX-2 (denoted 5.8) (Holtman *et al.*, 1996, *supra*). LOX-1
- 20 immunoprecipitation from extracts of cv Vintage leaf tips removed 85% of (LOX-1) protein and 15% of LOX-2 protein.

- Immunoprecipitation was performed in a V-bottom 96-well plate by adding 5  $\mu$ l 5D2-coated Dynabeads (Dynal) and 75  $\mu$ l buffer [20 mM Tris-HCl pH 7.5, 1% v/v Bovine Calf Serum (HyClone)] to 20  $\mu$ l of each
- 25 leaf tip extract. The plate was incubated on a titerplate shaker (MTS4, IKA, Labor Technik) for 1 hour at 4°C. The immunoprecipitate was pelleted by centrifugation at 4°C in a Sigma 302-K centrifuge for 10 minutes at 2000 rpm. The supernatant (70  $\mu$ l) from each sample was assayed for lipoxygenase activity in a flat bottom 96 well plate, as
- 30 described below, but with addition of 100  $\mu$ l assay buffer (25 mM HEPES, 0.2 M boric acid, pH 6.5).

The LOX-1 and LOX-2 assays were adapted for a high-throughput screening method. Leaf tips (1 cm) from eight 4 day-germinated grains were individually homogenised in 150 µl ice-cold buffer (20 mM Tris-HCl, pH 7.5) for 2 x 30 seconds in a multi-well homogeniser (Berg *et al.*, 1992, *Electrophoresis* 13: 76-81). After centrifugation for 15 minutes at 3000 rpm, 40 µl of the supernatant of each extract was transferred to a flat bottom 96 well plate. To each well, 170 µl buffer (25 mM HEPES, 0.2 M boric acid pH 6.5,  $1 \cdot 10^{-5}$  M NDGA) and 10 µl substrate (20 mM linoleic acid) were added and then incubated for 20 minutes at 25°C. The reaction was terminated by the addition of 20 µl saturated potassium iodide solution (KI) and incubated for a further 8 minutes at 25°C. The redox reaction between hydroperoxydienes and KI yields  $I_2$ , which was monitored by its extinction maximum at 350 nm in a microplate reader (Multiskan MCC/340).

15

### 3. Identification of potential lipoxygenase 1 mutants in the M3 and M4 grain of mutagenised barley

Grain of the M3 generation of cv Vintage and cv Caruso was stored at 45°C for 6.5 days to break dormancy, ensuring a 95% germination frequency. M3 grain of cv Vintage (9318) and cv Caruso (9633) was germinated and screened for lines whose LOX-1 activity was 15% or less of wild-type grain. The putative mutant lines (50 cv Vintage and 42 cv Caruso lines) were propagated to the M4 generation, harvested, and the germinated grain re-screened. The mutant LOX-1 phenotype was confirmed in one cv Vintage line and six cv Caruso lines, after measuring the lipoxygenase activity in extracts of 5 leaf-tips from each line. When the LOX-1 and LOX-2 activities in germinating embryos of these 7 putative mutants were examined, only the cv Vintage mutant (denoted Line G) showed a major reduction in LOX-1. In mature quiescent grain, lipoxygenase activity present in the embryo is almost exclusively LOX-1 activity, due to the differential expression pattern of the two isoenzymes (Schmitt and van Mechelen, 1997, *Plant Sci.* 128: 141-150). The total

30

lipoxygenase activity in extracts of embryos from Line G mature dry grain (M5 generation) was  $0.06 \pm 0.04$  U/mg protein in comparison to  $0.74 \pm 0.44$  U/mg protein in cv Vintage embryo extracts, as determined by the spectrophotometric lipoxygenase assay described in section 2 of Example

- 5 1. The residual lipoxygenase activity in mature embryos of Line G in both the M4 and M5 generations was found to be approximately 9 % of the parental line.

### Example 2

10 **Line G is a cv Vintage Mutant with a**  
**Low-Lipoxygenase Phenotype**

- The agronomic properties and mutant phenotype of Line G were analysed in material of the M5 generation. Initial analyses were
- 15 conducted to confirm that the analysed M5 material was homozygous for the mutant phenotype. The low LOX-1 phenotype in Line G, detected in the M3 generation, could result from a dominant or a recessive mutation. If the Line G selected at the M3 generation was heterozygous for a dominant mutation, then subsequent generations would show segregation
- 20 for the phenotype. The lipoxygenase activity in 26 individual Line G embryos from quiescent grain of the M5 generation was measured and compared to cv Vintage wild type embryos. The lipoxygenase activity in all Line G embryos was very low, with an average of  $0.06 \pm 0.04$  U
- 25 lipoxygenase per mg protein, compared to  $0.74 \pm 0.44$  U lipoxygenase per mg protein in wild type cv Vintage embryos. These data confirmed that Line G in the M5 generation was homozygous for the low lipoxygenase trait.



**1. Line G has a wild type plant growth physiology and grain development.**

Line G and cv Vintage grain were germinated and grown in a climate chamber under 16 hours light at 15°C and 8 hours dark at 12°C at a relative humidity of 80%. The growth characteristics of Line G and cv Vintage plants were similar with regard to plant height, number of tillers per plant, the onset of flowering and number of grains per spike. The fresh weight (**Figure 2**) and dry weight (**Figure 3**) of grain of Line G and wild type cv Vintage during development from 5 days after flowering (DAF) until full maturity, approximately 90 DAF, were very similar.

**2. Line G grain have a low-lipoxygenase 1 phenotype throughout development**

Lipoxygenase activity was measured in extracts of developing barley grain of Line G (M5 generation) and wild type cv Vintage. Grain was homogenised in ice-cold 20 mM Tris-HCl buffer pH 7.5 containing 0.1% (v/v) Nonidet P-40, a non-ionic detergent that enhances lipoxygenase extraction, and centrifuged at 15,000 *g* for 20 minutes to remove insoluble material. Lipoxygenase activity in the extracts was measured polarographically in 200 µl oxygen-saturated buffer (0.2 M boric acid, 25 mM HEPES, pH 6.5) containing 1.2 mM linoleic acid at 25°C, using a Clark-type electrode to measure oxygen consumption. Lipoxygenase activity increased during the first 20 days of grain development in both Line G and wild-type grain, but only in Line G did the activity level fall during grain maturation (**Figure 4**).

The relative amounts of 9-HPOD and 13-HPOD formed during linoleic acid oxidation provides a measure of the levels of LOX-1 and LOX-2 activity in the grain extracts. In this case Nonidet P-40 was omitted from the grain extraction buffer to avoid the co-extraction of hydroperoxide-consuming enzymes. The extracts (100 µl), mixed with 10 ml 50 mM phosphate buffer pH 6.5 containing 200 µM linoleic acid, were incubated for 20 minutes. The reaction was terminated by adjusting

the pH to 3.5, and an internal standard was added. The hydroperoxides formed in the assay were bound on an octadecyl solid phase column (Bakerbond, Baker) and eluted with methanol. The 9-HPOD and 13-HPOD were then separated by reverse phase HPLC on a C-18 column with an isocratic elution solvent (tetrahydrofuran:Methanol:H<sub>2</sub>O:acetic acid; 25:30:44.9:0.1 (v/v) adjusted to pH 5.5 with concentrated ammonia) at a flow rate of 0.5 ml/minute as described by Aarle *et al.*, 1991, *FEBS Letters* 280: 159-162. Hydroperoxides were detected at 234 nm and the HPOD peaks were corrected against the internal standard, prostaglandin B2.

**Figure 5** shows that 13-HPOD was the major product of lipoxygenase activity present in grain during the first 20 DAF, while 9-HPOD was formed by lipoxygenases active during grain maturation. While both Line G and wild-type grain extracts shared a similar profile of 13-HPOD synthesising activity, Line G did not show the wild-type rise in 9-HPOD synthesising activity. These data are consistent with a loss of LOX-1 activity in maturing Line G barley grain.

### 3. Line G grain have a low-lipoxygenase 1 phenotype on germination

Total lipoxygenase activity in extracts of embryos of grain germinated at 15°C was assayed as described in Example 1. The lipoxygenase activity present in quiescent wild-type grain declined during the first 4 days of germination and then increased (**Figure 6**). In Line G, lipoxygenase activity in quiescent grain was very low but increased after 4 days.

Analysis of the HPODs formed by the lipoxygenase activity in germinating embryos showed that 9-HPOD was the major product of lipoxygenases present in quiescent wild-type grain (**Figure 7**). The level of 9-HPOD formation fell with the decline in lipoxygenase activity in the extracts. The rise in lipoxygenase activity after 4 days was accompanied by the formation of both 9-HPOD and 13-HPOD. The low lipoxygenase

activity in Line G quiescent grain was associated with an absence of HPOD formation, while the rise in activity after 4 days mainly produced 13-HPOD. These data provide evidence that LOX-1 activity leading to the formation of 9-HPOD is greatly reduced in the embryos of both developing, quiescent and germinating barley grain of Line G, while LOX-2 activity leading to formation of 13-HPOD is unchanged in Line G.

### Example 3

#### **Line G has a Mutant Lipxygenase 1 Gene (*lox-1*) Causing a Low Lipxygenase Phenotype**

The molecular basis for the low-LOX-1 phenotype of Line G was investigated in order to provide a complete description of the mutant. The following analyses were performed to provide a complete characterization of the phenotype:

##### **1. Lipxygenase-1 is synthesised in the developing and germinating grain of Line G**

Western blot analysis of extracts of embryos from developing and germinating barley grain were performed in parallel with the measurement of lipxygenase activity, as described in Example 2. The crude extracts were separated by sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) according to Laemmli, 1970, *Nature* 227: 680-685. The separated proteins were transferred to nitrocellulose by semi-dry blotting, according to Towbin *et al.*, (1979) *Proc. Natl. Acad. Sci. USA* 76: 4350-4354. The blot was probed with the LOX-1 specific monoclonal antibody, 5D2, as described Holtman *et al.*, 1996, *Plant Physiology* 111: 569-576, at 500x dilution, followed by incubation with goat anti-mouse antibody coupled to alkaline phosphatase, and detected with the alkaline phosphatase substrates nitro blue tetrazolium and 5-bromo-4-chloro-3-indolyl-phosphate as described

by Holtman *et al.*, 1996, *Plant Physiol.* 111: 569-576. The Western analyses revealed that LOX-1 protein was detected in developing grain from 10 DAF in cv Vintage embryos and the level increased during grain maturation (**Figure 8**). The protein was also present in the embryo of cv  
5 Vintage quiescent grain but declined slowly during germination (**Figure 9**). Although LOX-1 is recognised in extracts of Line G embryos and migrates in SDS-PAGE as a protein of similar size to cv Vintage LOX-1, the immunodetectable levels of the protein in Line G were slightly lower than in cv Vintage.

10

## 2. The *lox-1* gene is expressed in the developing and germinating grain of Line G

Total RNA was isolated from embryos of developing and germinating barley grain, according to the procedure of Hensgens and  
15 van Os-Ruygrok, 1989, *Rice Genet. Newslett.* 6: 163-168, in parallel with the measurement of lipoxygenase activity, described in Example 2. The RNA samples (7.5 µg) were separated on denaturing agarose gels and Northern blotted as described by Sambrook *et al.*, 1989 in Molecular Cloning, a Laboratory Manual, Cold Spring Harbour Laboratory Press,  
20 NY. The blots were hybridised with a <sup>32</sup>P-labelled probe generated from the barley 3' untranslated region, nucleotides 2659-2801 [SEQ ID NO:1], of the *lox 1* cDNA (EMBL Accession no. L35931) as described by Holtman *et al.*, 1996 *Plant Physiol.* 111: 569-576, using the Amersham Random Prime Kit.

25 *Lox-1* transcripts encoding LOX-1 were detected in embryos of developing and mature cv Vintage and Line G grain from 30 DAF (**Figure 10**). The level of *lox-1* transcripts increased during germination in both cv Vintage and Line G embryos, indicating *de novo* expression of the *lox-1* gene (**Figure 11**). Since the detectable levels of *lox-1*  
30 transcripts were similar in Line G and cv Vintage embryos, neither reduced *lox-1* transcription or transcript stability can account for the low-lipoxygenase phenotype of Line G.

### 3. The *lox-1* gene of Line G encodes a mutant form of lipoxygenase-1

The nucleotide sequence of the *lox-1* gene of Line G and cv Vintage were analysed and compared in order to determine the molecular basis for the low-LOX-1 phenotype of Line G, which is characterised by normal transcription of the *lox-1* gene, but reduced accumulation and activity in the expressed lipoxygenase enzyme in grain.

Genomic DNA from Line G and wild-type cv Vintage was isolated from seedling leaf tissue according to a method described by Pich and Schubert 1993, *Nucleic Acids Res.* 21: 3328. The *lox-1* gene in the genomic DNA preparations was amplified by polymerase chain reaction (PCR) using primers based on the sequence of the barley *lox-1* gene (van Mechelen *et al.* 1995, *BBA* 1254: 221-225; Rouster *et al.*, 1997, *Plant J.* 11: 513-523). The position and sequence of the oligonucleotide primers used to amplify the *lox-1* promoter and coding regions, indicated in **Figure 12** were as follows:

Forward primer 5'-GAA AAG CTT GGA GGT AGA CGC TGC-3' [SEQ ID NO:2] and reverse primer 5'-TAT AGG ATC CTT GTT CTT GGC CTC CTC TCC TCG-3' [SEQ ID NO:3] were used to PCR amplify the *lox-1* promoter domain (-361 to +68) of Line G and cv Vintage *lox-1*.

Forward primer 5'-AGT GAA AAA CAG TGT GCT GGT G-3' [SEQ ID NO:4] and reverse primer 5'-GGC TTA AAG AGC AAC TGC TGA-3' [SEQ ID NO:5] were used to PCR amplify the Line G *lox-1* coding region.

Forward primer 5'-CAA GAT GCA TAT GCT GCT GGG AG-3' [SEQ ID NO:6] and reverse primer 5'-CGA TGG TTT AAA TTA GAT GGA GAT GCT GT-3' [SEQ ID NO:7] PCR amplified the cv Vintage *lox-1* coding region.

The PCR reactions consisted of 250 ng genomic DNA in a 50  $\mu$ l volume containing 50 pmol primer and 2 U *Pfu* DNA polymerase (Promega) according to the enzyme suppliers instructions. The PCR

amplifications were carried out in a Stratagene Robocycler: 1 minute at 94°C, 1 cycle; 1 minute at 94°C, 2 minutes at 62°C, and 5 minutes at 72°C, 30 cycles; 10 minutes at 72°C, 1 cycle. The PCR products were separated on 1.2 % agarose gels. DNA fragments, corresponding in length to the amplified region, were purified using Qiax II Gel extraction kit (Qiagen) and cloned into the plasmid pcDNA2.1 (Invitrogen). The nucleotide sequence of both strands of the cloned *lox-1* promoter and coding regions was determined using the dideoxynucleotide chain termination reaction with specific oligonucleotide primers and analysed on an ABI PRISM® 310 Genetic Analyzer (PE Biosystems). Sequence comparisons were performed using the DNA STAR sequence analysis software package (DNA STAR Inc., USA).

The promoter region and intron-exon structure of the barley *lox-1* coding region are shown in **Figure 13**, and were deduced from a comparison of the nucleotide sequence of the wild-type *lox-1* genomic and cDNA sequences (**Figure 12**). The sequenced region of the *lox-1* promoter region from -363 to +68, (numbered relative to the determined transcription start site; van Mechelen *et al.*, 1995, *BBA* 1254: 221-225), is sufficient to direct embryo-specific and temporally-regulated gene expression characteristic of the native gene (Rouster *et al.*, 1998, *Plant J* 15: 435-440). The promoter and transcribed region of the wild-type *lox-1* gene [SEQ ID NO:8] is 4663 nt in length and contains 6 introns of between 82 nt and 634 nt in length, which are absent from the respective cDNA [SEQ ID NO:10] and must therefore be removed during RNA transcript splicing.

Comparison of the nucleotide sequence of *lox-1* of Line G with that of wild-type (**Figure 12**) showed that the Line G *lox-1* allele has two point mutations. One is a silent C→T substitution at position 221 in exon 1, and the second is a G→A substitution at position 2347 in exon 3 (**Figure 13**). The wild-type barley *lox-1* gene encodes a protein of 862 amino acid residues [SEQ ID NO:9], while the mutation at position 2347

in Line G *lox-1* allele causes an amino acid substitution of glycine to aspartic acid at residue 368 in the encoded protein.

Alignment of related plant lipoxygenases indicated that the glycine-368 in barley LOX-1, is strongly conserved. Furthermore this  
5 residue, which corresponds to glycine-353 in soybean LOX-1, is one of 51 neutral or hydrophobic residues which line the substrate cavity of the enzyme, as seen from its crystal structure (Minor *et al.*, 1996, *Biochemistry* 35: 10687-10701) and is shown in (Figure 22). The insertion of a charged amino acid residue at this position is thus likely to  
10 disturb the structural and functional properties of the enzyme.

**4. The mutated LOX-1 protein encoded by the Line G *lox-1* allele has low enzymic activity and is responsible for the low lipoxygenase phenotype of Line G.**

15 The sodium azide mutagenesis of cv Vintage grain, which induced the mutated *lox-1* allele in Line G, may have induced additional mutations in the Line G genome. Two experimental approaches have been taken to demonstrate that the mutant *lox-1* allele in Line G is responsible for its low lipoxygenase phenotype, rather than other  
20 mutations in the genome. The enzymic activity of the LOX-1 encoded by the mutant and wild-type *lox-1* allele have been determined in order to prove that the glycine→aspartic acid substitution in the mutant enzyme causes reduced stability and activity. The two *lox-1* genes were transiently expressed in aleurone protoplasts isolated from imbibed  
25 mature grain, since the level of endogenous lipoxygenase expression in these cells was expected to be below detection limits. None of the identified barley lipoxygenase genes, which are expressed in germinating barley, are detected in the aleurone tissue (van Mechelen *et al.*, 1999 *supra*). In order to direct transient expression of the *lox-1* gene in  
30 aleurone protoplasts, their coding regions were translationally fused to a constitutive promoter known to be active in these protoplasts.

The coding regions of the mutant (sequence positions +1 to +4350) and the wild-type *lox-1* gene (sequence positions +69 to +4230) and the wild-type *lox-1* cDNA (sequence positions +69 to +2654) each cloned in plasmid pcDNA2.1 (see section 3), were excised by digestion  
5 of the *KpnI* and *EcoRV* sites in vector polylinker. The coding regions were cloned in the pUBARN plasmid (Jensen *et al.*, 1998, *Hereditas* 129: 215-225) between the constitutively active maize ubiquitin *Ubi* promoter (as described in U.S. patent No. 005510474A) and the *Nos* terminator, in place of the *bar* gene which encodes phosphinotricin acetyl transferase  
10 (Figure 14).

Protoplasts were isolated from aleurone tissue of imbibed *Hordeum vulgare* cv Himalaya according to the protocol of Skriver *et al.* 1991, *Proc. Natl. Acad. Sci. USA* 88: 7266-7270. Aliquots of  $2 \cdot 10^5$  protoplasts were transfected at 0°C with ~100 µg plasmid DNA  
15 (equimolar amounts of each plasmid) by polyethylene glycol (PEG) mediated DNA uptake (Lee *et al.*, 1997, *Plant Mol. Biol.* 13: 21-29), and then incubated in aleurone protoplast culture media at 25°C as described previously (Skriver *et al.*, 1991 *supra*). After 48 hours incubation, the culture medium was carefully removed and the protoplasts were re-  
20 suspended and homogenised in 300 µl lipoxxygenase assay buffer (0.2 mM boric acid, 25 mM HEPES, pH 6.5). The homogenates were centrifuged at 15,000 *g* for 5 minutes to pellet insoluble material, and the supernatants (10 µl) were subsequently assayed for total lipoxxygenase activity using the rapid screening assay described in Example 1, section  
25 1, but with omission of the NDGA inhibitor. The protein content of the protoplast extracts was measured by a Bradford dye-binding assay (Bradford 1976, *Anal. Biochem.*, 72: 248) supplied by Bio-Rad Laboratories, Hercules, California, USA, and lipoxxygenase activity was expressed per mg protein in the extract.

30 Protoplasts transfected with the control plasmid, pUBI-GUS, where the maize ubiquitin-1 promoter directs expression of the  $\beta$ -glucuronidase reporter gene, gave no detectable lipoxxygenase activity.



Transient expression of the wildtype *lox-1* gene and cDNA in protoplasts both gave high levels of lipoxygenase activity in the protoplast extracts (Figure 15). The higher expression of the wild-type *lox-1* cDNA in comparison to the genomic sequence may be due to a higher transfection frequency for the smaller *lox-1* cDNA expression plasmid (4929 bp versus the 6505 bp *lox-1* gene construct). Transient expression of the mutant *lox-1* gene gave low levels of lipoxygenase activity, ~10 % of wild-type lipoxygenase activity. These data clearly demonstrate that the mutant *lox-1* gene in Line G encodes a lipoxygenase with greatly reduced activity, which accounts for the low lipoxygenase phenotype.

#### Example 4

##### PCR-Cleavage Amplified Polymorphic Site (PCR-CAPS) Assay:

##### A Method Used for Identification of the Mutant *lox-1* Gene

15

An analytical method allowing the identification of the Line G mutant *lox-1* gene in any genetic background was developed based on the PCR-CAPS assay. The assay involves PCR amplification of genomic DNA fragments, followed by digestion of the amplified sequences with a specific restriction endonuclease to display a restriction fragment length polymorphism (RFLP).

The coding sequence of the mutant *lox-1* gene harbours two point mutations (see Example 3), where the mutation at position 2347 (Figure 12) introduces an additional *Aat* II restriction endonuclease cleavage site, not found in the wild-type *lox-1* gene (Figure 16). The following PCR-CAPS assay, based on the polymorphism created by the presence of this restriction site in the *lox-1* gene, is shown to discriminate between a wild-type *lox-1* gene and a mutated *lox-1* gene.

Genomic DNA was isolated from young leaves of M6 seedlings of *Hordeum vulgare*, L. cv Vintage and Line G according to the procedure of Pich and Schubert (1993, *supra*). The DNA sequence encompassing position 2347 (Line G *lox-1* gene mutation site) was

amplified by PCR, using primers specific for the *lox-1* gene [SEQ ID NO:8]. The DNA fragments amplified by the selected forward primer 5'-CGCTACGACGTCTACAACGA-3' [SEQ ID NO:13] and reverse primer 5'-CAGACTACTTTTGGCGGGA-3' [SEQ ID NO:14] are  
5 shown in **Figure 17**. PCR reactions were carried out with 250 ng genomic DNA in a 50- $\mu$ l volume containing 50 pmol of each primer and 1 unit Taq DNA polymerase (Promega) according to the suppliers instructions. PCR amplifications were carried out on a Stratagene Robocycler as follows: 1 minute at 94°C, 1 cycle; 1 minute at 94°C, 1.5  
10 minutes at 60°C, and 2 minutes at 72°C, 30 cycles; 10 minutes at 72°C, 1 cycle. The amplified fragments of the mutant and wild-type *lox-1* gene were ~650 bp (**Figure 18**), corresponding to the expected size (**Figure 17**). The PCR products, purified on a spin column (Qiagen), were digested with 25 unit *Aat* II restriction endonuclease for 24 hours at 37°C  
15 and analyzed on a 1.2% agarose gel.

Digestion of the wild-type *lox-1* PCR product yielded DNA fragments of 10, 179, and 462 bp, and the fragments from the mutant *lox-1* PCR product were 10, 149, 179, and 313 bp, where additional DNA fragments were due to partial digestion of the *lox-1* PCR product (**Figure**  
20 **19**). The fragment pattern corresponds to the expected RFLP resulting from this mutation, where the 313 bp fragment is unique to the mutant *lox-1*. This PCR-CAPS assay provides a reproducible and specific tool for identification of the *lox-1* allele in barley and can thus be exploited in barley breeding programs aimed at introducing this gene in new barley  
25 varieties.

### Example 5

#### **Back-Crossing the Low Lipxygenase Phenotype of Line G to cv Alexis Demonstrates a Genetic Linkage to the Mutant *lox-1* Gene**

30 Repeated back-crossing was used to transfer the low-lipxygenase phenotype from line G into a recurrent parent (in this case the cv. Alexis). The back-crossing program shown in **Figure 20**, combined with selection

for the low-lipoxygenase phenotype, progressively substitutes the Line G genome by the recurrent parent genome. Furthermore, other mutations introduced into the Line G genome during the sodium azide mutagenesis treatment will be eliminated. In the first back-cross of the homozygous  
5 low-lipoxygenase Line G (denoted genotype *ll*) to cv Alexis (denoted genotype *LL*) the progeny lines will be heterozygous (denoted genotype *Ll*). A low-lipoxygenase phenotype due to a recessive mutation will not be detectable in lines heterozygous for the mutation. The progeny are self-pollinated and will give a normal Mendelian segregating population,  
10 namely 1 *LL* : 2 *Ll* : 1 *ll*. The low lox homozygous *ll* genotype resulting from the first back-cross will have 50 % cv Alexis genetic background. After ten rounds of back-crossing, the recurrent parent background will be approximately 99.9 %.

*Hordeum vulgare*, L. cv Alexis and Line G were propagated in a  
15 greenhouse throughout the back-crossing program. Back-crossed progeny grains were germinated in petri dishes on filterpaper, soaked with 4 ml H<sub>2</sub>O, for 3 days at 22°C in the dark. The low-lipoxygenase lines were screened by measuring total lipoxygenase activity in extracts of the coleoptile (top 7 mm) from the germinating seedlings, as described in  
20 Example 1. Progeny of the 3<sup>rd</sup> and 4<sup>th</sup> back-cross were also analysed for inheritance of the mutant *lox-1* gene using the PCR-CAPS assay described in Example 4.

The expected frequency of the low-lipoxygenase phenotype in the segregating progeny of the four back-cross generations was 25% for a  
25 recessive mutation. The observed frequency of low-lipoxygenase activity in the progeny (24 grains) of the four back-cross generations is in agreement with the expected frequency (Figure 20). When the 3<sup>rd</sup> and 4<sup>th</sup> back-cross progeny having the low lox homozygous *ll* genotype were analysed with the PCR-CAPS assay, they were all found to have the  
30 diagnostic 313 bp fragment, while progeny having wild-type lipoxygenase activity lacked this fragment (Figure 21).

The back-crossing program demonstrates that the mutant *lox-1* allele can be transferred to a new genetic background and is inherited in a recessive monofactorial manner following Mendelian segregation. Since the recurrent parent background is 93.8% in the 4<sup>th</sup> back-cross progeny, the co-inheritance of the mutant *lox-1* gene and the low-lipoxygenase phenotype provides confirmation of their genetic linkage.

#### Example 6

#### **Beer Brewed From Line G Barley Malt Accumulates Less *trans*-2-nonenal During Storage, Giving an Improved Flavour Stability**

*Hordeum vulgare* L cv Vintage and Line G were propagated in the field over several seasons in order to provide sufficient grain for industrial malting. The following industrial scale malting and brewing trials as well as analyses of the finished beer were performed to demonstrate the value of the Line G low-lipoxygenase barley for improved flavour stability.

##### **1. Industrial malting and kilning of Line G and cv Vintage**

Malting was performed on a 10-ton scale in an industrial malthouse in two trials as follows:

##### **Trial 1: Line G barley grain (1996 harvest)**

Steeping conditions: 8 hours wet; 14 hours dry; 8 hours wet; 10 hours dry; 4 hours wet in 16°C steeping water. Malting conditions: 12 hours at 18°C; 24 hours at 16°C; 24 hours at 14°C; 60 hours at 12°C.

Kilning conditions: 12 hours at 60°C; 3 hours at 68°C; 4 hours at 74°C; 3 hours at 80°C.

##### **Trial 2: cv Vintage and Line G (1996/1997 harvest)**

Steeping conditions: 8 hours wet; 10 hours dry; 6 hours wet; 15 hours dry; 4 hours wet in 15°C steeping water. Malting conditions: 5 days with inlet air at 15°C and spraying to maintain moisture level.

Kilning conditions: 10 hours at 50°C; 2 hours at 60°C; 2.5 hours at 80°C.

Malting analyses of 2 samples of the Line G malt from Trial 1 compared to the control malt, cv Nevada (Table 1) and from Trial 2 compared to cv Vintage (Table 2) confirmed that Line G malt was suitable for brewing trials.

TABLE 1

MALTING TRIAL 1				
Barley variety		cv Nevada	LineG	LineG
Crop year		1996	1996	1996
Malt analyses				
Moisture content	%	4.7	4.3	4.4
Extract fine as is.	%ai.	76.9	76.1	75.3
Extract fine d.m.	%dm.	81.4	79.5	78.7
Saccharification time	Min	< 10	< 10	< 10
Diastatic power	%WK	252	373	365
Color	EBC	2.8	4.4	3.8
pH		6.16	5.97	5.99
Turbidity	EBC	9.0	2.5	2.4
Total protein d.m.	%	10.35	10.74	12.03
Soluble nitrogen	mg/l	647	787	765
Sol. Protein % malt d.m.	%dm.	3.7	4.4	4.3
Kolbach		35.3	40.8	35.4
Free Amino Nitrogen	mg/l	97	125	118
Friability	%	89.5	85.6	89.5
Whole unmodified grains	%	1.1	0.6	0.5
Partly unmodified grains	%	2.3	1.0	0.6
$\beta$ -glucan in wort	mg/l	114	66	36
$\beta$ -glucan in malt	% w/w	0.24	0.11	0.05
S-methylmethionine/DMS eq.	$\mu$ g/g	2.4	6.4	8.4
Free DMS	$\mu$ g/g	1.0	6.6	4.7
NDMA	$\mu$ g/kg	n.d.	0.3/0.6	0.3/0.3

TABLE 2

MALTING TRIAL 2				
Barley variety		Vintage	Line G	Line G
Crop year		1996	1996	1997
Moisture content	%	4.1	4.1	4.3
Extract fine as is.	%ai.	77.0	75.6	77.5
Extract fine d.m.	%dm.	80.3	78.8	80.9
Fine/coarse difference	% dm.	0.7	1.6	1.7
Saccharification time	min	-	-	< 10
Diastatic power	%WK	343	342	268
Color	EBC	2.5	2.8	3.4
PH		6.05	6.01	6.12
Turbidity	EBC	1.5	1.3	2.5
Total protein d.m.	%	10.98	12.22	9.82
Soluble nitrogen	mg/l	696	741	610
Sol. Protein % malt d.m.	%dm.	3.9	4.1	3.4
Kolbach		35.2	33.7	34.6
Free Amino Nitrogen	mg/l	110	117	100
Friability	%	91.3	81.8	89.5
Whole unmodified grains	%	0.7	1.1	1.3
Partly unmodified grains	%	1.0	2.7	2.7
$\beta$ -glucan in wort	mg/l	97	172	117
Beta-glucan in malt	% w/w	-	-	0.3
S-methylmethionine/DMS eq.	$\mu$ g/g	-	-	-
Free DMS	$\mu$ g/g	-		-

**2. Industrial brewing with Line G, cv Vintage malt, and control malt cv Nevada**

Two brewing trials were performed, using wort prepared from Line G and control malt cv Nevada malt in Trial 1 and from Line G and control malt cv Vintage in Trial 2.

Beer was brewed on a 30-hl industrial scale with 475 kg malt according to the following scheme: Mashing in at 50°C; 30 minutes at 50°C; 30 minutes heating from 50 - 70°C; 15 minutes at 70°C. A portion of the wort was heated for 20 minutes from 70 - 100°C and 5 minutes at 100°C, while the main mash was kept at 70°C for another 25 minutes and then the two mashes was combined and kept for 10 minutes at 76°C. The brewing steps of wort boiling, whirlpool separation of spent grain, cooling, fermentation, lagering and packaging in brown glass bottles were according to standard brewing practise.

### **3. Flavor stability and T2N content of beer brewed from Line G, cv Vintage malt and control malt cv Nevada**

The freshly bottled beer was stored at 5°C and analysed within 2 months of production. The flavor-stability of the fresh and stored beer was evaluated in two independent laboratories following two different types of beer storage conditions. In laboratory A the beer was subjected to a forced aging process, where the beer was stored at 37°C for a period of 7 days, while in laboratory B the beer was stored at 30°C for 6 and 12 weeks. *Trans*-2-nonenal levels in beer were determined by gas chromatography and mass spectrometric detection following derivatisation of carbonyls with O-(2,3,4,5,6-pentafluorobenzyl)-hydroxylamine, essentially as described by Grönqvist et al. 1993 *Proceedings of the 24<sup>th</sup> EBC Congress*, Oslo, 421-428. A trained beer taste panel evaluated the overall flavor score of the beer, which includes detection of a cardboard flavor, indicative of free *trans*-2-nonenal in the beer.

#### **30 Laboratory A: Forced-Aging Tolerance**

Comparison of beer, brewed from Line G and the control malt, cv Nevada, in the first brewing trial (Table 3) demonstrated that beer from



Line G had a greater flavor stability and a lower *trans*-2-nonenal content following forced-aging as compared to the controls. The second trial, comparing beer brewed from Line G malt with beer brewed from cv Vintage malt, the parental cultivar, confirmed the initial data (Table 4).

5

TABLE 3

BREWING TRIAL 1		
Barley Malt	cv Nevada	Line G
SO <sub>2</sub> Content (mg/ml)	1	1
T2N** (ppb) – Fresh Beer	0.009	0.005
T2N** (ppb) – Aged Beer (37°C / 7 DAY)	0.117	0.025
Flavor* - Fresh Beer	5.9	5.3
Flavor* – Aged Beer (37°C / 7 DAY)	1.3	5.1

\* Flavor evaluation scale 1-10 of increasing quality; \*\**trans*-2-nonenal.

TABLE 4

BREWING TRIAL 2		
Barley Malt	cv Vintage	Line G
SO <sub>2</sub> Content (mg/ml)	2	2.5
T2N** (ppb) – Fresh Beer	0.023	0.019
T2N** (ppb) – Aged Beer (37°C / 7 Day)	0.078	0.035
FLAVOR* - Fresh Beer	5.5	6.1
FLAVOR* - Aged Beer (37°C / 7 Day)	2.9	5.9

\* Flavor evaluation scale 1-10 of increasing quality; \*\**trans*-2-nonenal.

10

#### Laboratory B: 30°C Storage Tolerance

Beer brewed from Line G malt had lower *trans*-2-nonenal levels following 6 and 12 weeks at the elevated storage temperature of 30°C, when compared to beer brewed from either of the reference malts (Table 5 and 6) and had a better flavor-stability as judged by a taste panel. The taste-threshold for *trans*-2-nonenal in these analysed beers lies close to 0.08 ppb.

15

TABLE 5

BREWING TRIAL 1		
Barley Malt	cv Nevada	Line G
<i>trans</i> -2-nonenal (ppb) – Fresh Beer	0.050	0.044
<i>trans</i> -2-nonenal (ppb) – 30°C / 6 weeks	0.072	0.037
<i>trans</i> -2-nonenal (ppb) – 30°C / 12 weeks	0.095	0.046
<i>trans</i> -2-nonenal flavor* – Fresh Beer	0.6	0.3
<i>trans</i> -2-nonenal flavor* – 30°C / 6 weeks	3.7	1.4
<i>trans</i> -2-nonenal flavor* – 30°C / 12 weeks	2.5	0.6

\* *trans*-2-nonenal flavor detection score on a scale of 1 -10

5

TABLE 6

BREWING TRIAL 2		
Barley Malt	cv Vintage	Line G
<i>trans</i> -2-nonenal (ppb) – Fresh Beer	0.070	0.062
<i>trans</i> -2-nonenal (ppb) – 30°C / 6 weeks	0.093	0.070
<i>trans</i> -2-nonenal (ppb) – 30°C / 12 weeks	0.133	0.080
<i>trans</i> -2-nonenal flavor* – Fresh Beer	0.3	0.9
<i>trans</i> -2-nonenal flavor* – 30°C / 6 weeks	2.5	1.7
<i>trans</i> -2-nonenal flavor* – 30°C / 12 weeks	2.2	1.3

\* *trans*-2-nonenal flavor detection score on a scale of 1 -10

- The improved flavor-stability of beer brewed from Line G malt, as measured by the levels of *trans*-2-nonenal detected in the beer
- 10 following storage at 30°C from the combined brewing trial data, is shown to be statistically significant (Table 7).

**TABLE 7**  
**TRANS-2-NONENAL IN STORED BEER**

<b>fresh</b>	<b>Mean</b>	<b>Stdev</b>	<b>Difference</b>	<b>2-tailed p</b>	<b>Significant (p &lt; 0.05)</b>
reference	0.060	0.012	0.007	0.34	no
line-G	0.053	0.011			

<b>6 weeks, 30 °C</b>	<b>Mean</b>	<b>Stdev</b>	<b>Difference</b>	<b>2-tailed p</b>	<b>Significant (p &lt; 0.05)</b>
reference	0.083	0.013	0.029	0.031	yes
line-G	0.054	0.021			

5

<b>12 weeks, 30 °C</b>	<b>Mean</b>	<b>Stdev</b>	<b>Difference</b>	<b>2-tailed p</b>	<b>Significant (p &lt; 0.05)</b>
reference	0.114	0.023	0.051	0.003	yes
line-G	0.063	0.020			

Since the natural sulfite levels were low in both brewing trials, the free *trans*-2-nonenal levels in the aged beer would closely reflect the *trans*-2-nonenal potential of the different beers, namely the level of *trans*-2-nonenal adducts present in the fresh beer. Addition of sulfite can temporarily delay the staling process, by complexing free-*trans*-2-nonenal, until sulfite levels are reduced by oxidation due to gaseous exchange through the packaging.

The described brewing trials with low-LOX-1 barley malt provide the first unequivocal evidence that LOX-1 activity in barley during the malting and brewing process is a key determinant of the appearance of the off-flavor compound *trans*-2-nonenal in aged beer.

The above specification includes citations to numerous publications. Each publication is hereby incorporated by reference for all purposes, as if fully set forth.

Applicant's or agent's file reference 11225.11W001	International application No.
---	-------------------------------

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM  
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page <u>16</u> , line <u>6-10</u>	
B. IDENTIFICATION OF DEPOSIT <span style="float: right;">Further deposits are identified on an additional sheet <input type="checkbox"/></span>	
Name of depositary institution National Collections of Industrial, Food and Marine Bacteria (NCIMB)	
Address of depositary institution (including postal code and country) 23 St. Machar Drive, Aberdeen, AB243RY, Scotland, United Kingdom	
Date of deposit 19 December 2000	Accession Number (Unknown)
C. ADDITIONAL INDICATIONS (leave blank if not applicable) <span style="float: right;">This information is continued on an additional sheet <input type="checkbox"/></span>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

<p align="center">For receiving Office use only</p> <p><input checked="" type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer A. Groeg.</p>	<p align="center">For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p>
---	---

## WE CLAIM:

1. A barley plant or portion thereof comprising a mutant LOX-1 protein, the plant or plant portion characterized by a reduction or absence of lipoxygenase activity as compared to a non-mutated control.
2. The barley plant or portion thereof of claim 1, wherein the lipoxygenase activity comprises catalysis of oxidation of free and esterified polyunsaturated fatty acids and polyunsaturated octadecanoic fatty acids to form 9-hydroperoxy fatty acid derivatives.
3. The barley plant or portion thereof of claim 1, wherein the LOX-1 protein is encoded by a mutated *lox-1* nucleic acid sequence having one or more mutation.
4. The barley plant or portion of claim 3, wherein said one or more mutation is induced by chemical mutagenesis or radiation.
5. The barley plant or portion of claim 3, wherein said one or more mutation is induced by site-directed mutagenesis.
6. The barley plant or portion thereof of claim 3, wherein said one or more mutation is positioned within the promoter or transcribed region of the *lox-1* nucleic acid sequence.
7. The barley plant or portion of claim 1, wherein said mutant LOX-1 protein comprises an acidic, basic, or polar amino acid substitution at one or more conserved neutral or hydrophobic amino acid residue lining the substrate cavity of wild-type barley LOX-1.

8. The barley plant or portion of claim 7, wherein said mutant LOX-1 protein comprises the amino acid sequence of SEQ ID NO: 12, wherein Xaa is an acidic, basic, or polar amino acid.
9. The barley plant or portion of claim 8, wherein Xaa is glutamic acid or aspartic acid.
10. The barley plant or portion of claim 9, wherein Xaa is aspartic acid.
11. A transgenic barley plant or portion thereof comprising a heterologous nucleic acid sequence expressing an antisense sequence to at least a portion of a transcribed region of the barley *lox-1* gene, the heterologous sequence operably linked to a gene promoter sequence and a transcription terminator sequence.
12. The transgenic barley plant or portion of claim 11, wherein said gene promoter directs constitutive expression of the antisense sequence.
13. The transgenic barley plant or portion of claim 11, wherein said gene promoter directs tissue-specific or temporally-regulated expression of the operably linked antisense sequence in tissues of the plant.
14. The transgenic barley plant or portion of claim 11, wherein said tissue is developing, germinating, or mature embryonic tissue.
15. The transgenic barley plant or portion of claim 11, wherein said gene promoter sequence is a barley *lox-1* gene promoter or a portion thereof.
16. Grain or plant progeny produced from the barley plant or plant portion of any of claims 1-15.

17. A plant product produced from the barley plant or plant portion of any of claims 1-15 or from the grain or plant progeny of claim 16.
18. The plant product of claim 17, wherein the product is malt.
19. A beverage manufactured using the plant product of any of claims 17-18.
20. Beer manufactured using the plant product of any of claims 17-18.
21. The use of a plant product according to any of claims 17-18 for the preparation of a beverage presenting organoleptic qualities that remain stable over a measured period of time or at elevated storage temperatures as compared to a beverage prepared from a plant product of a non-mutated control.
22. Use of a barley plant or portion thereof, grain or plant progeny, or plant product according to any of claims 1-18 for the manufacture of a beverage.
23. Use of a barley plant or portion thereof, grain or plant progeny, or plant product according to any of claims 1-18 for the manufacture of a malt or beer.
24. Use of a barley plant or portion thereof, grain or plant progeny, or plant product according to any of claims 1-18, for the stabilization of organoleptic properties of a brewed product over a measured period of time as compared to a control brewed product produced using a non-mutated barley plant or portion thereof, grain or plant progeny, or plant product.
25. Use of a barley plant or portion thereof, grain or plant progeny, or plant product according to any of claims 1-18 for the manufacture of a brewed product having reduced levels of free *trans*-2-nonenal over a measured period of time or under conditions of elevated storage temperature, as

compared to a control brewed product produced using a non-mutated barley plant or portion thereof, grain or plant progeny, or plant product.

26. A beverage produced from a barley plant or portion thereof, grain or plant progeny, or plant product according to any of claims 1-18, the beverage having a reduced *trans*-2-nonenal content over a measured period of time or under conditions of elevated storage temperature as, compared to a control beverage produced from a non-mutated barley plant or portion thereof, grain or plant progeny, or plant product.
27. The beverage of claim 26, wherein the beverage is beer.



FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8






FIGURE 9




FIGURE 10





FIGURE 11



## FIGURE 12A

(-288) Promoter

LG CAGCCCCATGCATGCACATGCACATGCAGTGCAGCAAGCACCGCTCGATGGCGATCACCCGTCAAG -208  
 WT CAGCCCCATGCATGCACATGCACATGCAGTGCAGCAAGCACCGCTCGATGGCGATCACCCGTCAAG

LG GGACCGGAGCGGCCATGCCAAGCACGAGGAGGGCA CGTCACCGTCCGGCGGCAGCACGTTGGAGAGCACGTGCCCGTCCG -128  
 WT GGACCGGAGCGGCCATGCCAAGCACGAGGAGGGCACGTCACCGTCCGGCGGCAGCACGTTGGAGAGCACGTGCCCGTCCG

LG ATCCATCTCTCCAAAGCCGAGCGGCCACACACCCGGGACCGGACCCGGCCTATAAATTGCCCGGACCGAGCTGCAA -48  
 WT ATCCATCTCTCCAAAGCCGAGCGGCCACACACCCGGGACCGGACCCGGCCTATAAATTGCCCGGACCGAGCTGCAA

Transcription start site (+1)

LG GCAGCTCCTCACACACTCAGCGCAACACACATCCATCTTCACTGAAAGTGAAGAAACAGTGTGCTGGTGCCATTGGTTG 32  
 WT GCAGCTCCTCACACACTCAGCGCAACACACATCCATCTTCACTGAAAGTGAAGAAACAGTGTGCTGGTGCCATTGGTTG

Translation start site (+69) →

LG GAGCAGTGAAAGCGAGGAGAGGAGGCCAAGAACAAAGATGCTGTGGAGGGCTGATCGACACCCCTCACGGGGCGGAACAA 112  
 WT GAGCAGTGAAAGCGAGGAGAGGAGGCCAAGAACAAAGATGCTGTGGAGGGCTGATCGACACCCCTCACGGGGCGGAACAA

LG GAGCGCCCGGCTCAAGGGCACGGTGGTGTCTCATGCGGAAGAACGTTGCTGGACCTCAACGACTTCGGCGCCACCATCATCG 192  
 WT GAGCGCCCGGCTCAAGGGCACGGTGGTGTCTCATGCGGAAGAACGTTGCTGGACCTCAACGACTTCGGCGCCACCATCATCG

(Silent mutation)\*

LG ACGGCATCGGGGAGTTCCCTCGGCAAGGGTGTCACTGCCAGCTTATCAGCTCCACCGCCGTCGACCAAGGTAATCACTAC 272  
 WT ACGGCATCGGGGAGTTCCCTCGGCAAGGGGTCACCTGCCAGCTTATCAGCTCCACCGCCGTCGACCAAGGTAATCACTAC

LG CCTCCTCCGGCCTTCTCCTCTGTTTACAAGATATAGTATTTCTTTCTGTGGGCCGGCGCCATGGATGGATGGATGTGT 352  
 WT CCTCCTCCGGCCTTCTCCTCTGTTTACAAGATATAGTATTTCTTTCTGTGTGGGCCGGCGCCATGGATGGATGGATGTGT

## FIGURE 12B

LG CTGGATCGGCTAAAGAAGATAGGATAGCTAGCCCTGGCCGGTCGTCTTTACCTGAGCATGGGCATATGCCATCGAAAAAA 432  
 WT CTGGATCGGCTAAAGAAGATAGGATAGCTAGCCCTGGCCGGTCGTCTTTACCTGAGCATGGGCATATGCCATCGAAAAAA

LG GAGACAAACAGCATGCATGGTGGCGCACCAAGACCGAGAGCACCGGATGCTCGAGACAAAGCAACACAACAAGC 512  
 WT GAGACAAACAGCATGCATGGTGGCGCACCAAGACCGAGAGCACCGGATGCTCGAGACAAAGCAACACAACAAGC

LG AAGGACGACACGTCAAAAAGCAACAAAGCAAGGACGGCAGTCAAAAGCAACAAACCTAAAGCACAAAGA 592  
 WT AAGGACGACACGTCAAAAAGCAACAAAGCAAGGACGGCAGTCAAAAGCAACAAACCTAAAGCACAAAGA

LG CGTAAGAGCAAGCACACAATCAGCAGGCTATAAACAGTTGTTCATCAAAAAACAACGCTGGAAAGAGAGAGAAAGGAA 672  
 WT CGTAAGAGCAAGCACACAATCAGCAGGCTATAAACAGTTGTTCATCAAAAAACAACGCTGGAAAGAGAGAGAAAGGAA

LG GTAGTAGCCATGAAAAATTAAATCACCGGGCGTTGCTCTTTGCCCAACAATTAAATCAAGCAGGGTACGTGGCATGTATAG 752  
 WT GTAGTAGCCATGAAAAATTAAATCACCGGGCGTTGCTCTTTGCCCAACAATTAAATCAAGCAGGGTACGTGGCATGTATAG

LG TTCTTTGTAAGTAAACTAAGCATGTGATATAGAAAGGTACGTGGTGGTGAGACAAACGGCGGTCCGGGAAGGTGGCGCG 832  
 WT TTCTTTGTAAGTAAACTAAGCATGTGATATAGAAAGGTACGTGGTGGTGAGACAAACGGCGGTCCGGGAAGGTGGCGCG

LG GAGGCGGAGCTGGAGCAGTGGGTGACGAGCCTGCCGTGCTGACGACGGGGGAGTCCAAAGTTCGGCCTCACCTTCGACTG 912  
 WT GAGGCGGAGCTGGAGCAGTGGGTGACGAGCCTGCCGTGCTGACGACGGGGGAGTCCAAAGTTCGGCCTCACCTTCGACTG

LG GGAGGTGGAGAAGCTCGGGGTGCCGGGCGCCATCGTCGTCAACAACTACCACAGCTCCGAGTTCTTGCTTAAACCATCA 992  
 WT GGAGGTGGAGAAGCTCGGGGTGCCGGGCGCCATCGTCGTCAACAACTACCACAGCTCCGAGTTCTTGCTTAAACCATCA

LG CCCTCCACGACGTCCCCGGCCGAGCGGCAACCTCACCTTCGTGCCAACTCATGGATCTACCCCGCCGCAACTACCGA 1072  
 WT CCCTCCACGACGTCCCCGGCCGAGCGGCAACCTCACCTTCGTGCCAACTCATGGATCTACCCCGCCGCAACTACCGA

## FIGURE 12C

LG TACAGCCGCGTCTTCTTCGCCAACGACGTCGCTGGATTTTCCTCTACTTTTCCTCTCCTTTTCATTTTCACCGCCTTCGTCA 1152  
 WT TACAGCCGCGTCTTCTTCGCCAACGACGTCGCTGGATTTTCCTCTACTTTTCCTCTCCTTTTCATTTTCACCGCCTTCGTCA  
  
 LG TTCAATGGTCGATCATTAAGTCTTTGCCAGGACAAATAGATGATGAGCTAGGAGTGGTTACCACTTAGCAGTACGTACATTAT 1232  
 WT TTCAATGGTCGATCATTAAGTCTTTGCCAGGACAAATAGATGATGAGCTAGGAGTGGTTACCACTTAGCAGTACGTACATTAT  
  
 LG TTATTCCCGTGTGGTAGAAAAAGGATATGGTTTGGTGCAGATCGACACAAAGATTGAATGAAAAGTTGCACCGTGGCACCCGTG 1312  
 WT TTATTCCCGTGTGGTAGAAAAAGGATATGGTTTGGTGCAGATCGACACAAAGATTGAATGAAAAGTTGCACCGTGGCACCCGTG  
  
 LG GCAGCGTGGTAGGTGAAAAATAACTGTTGCACGGATCCACCCACATGATTGTTTTCATGAATAAACTTTTAAAGGATGTGT 1392  
 WT GCAGCGTGGTAGGTGAAAAATAACTGTTGCACGGATCCACCCACATGATTGTTTTCATGAATAAACTTTTAAAGGATGTGT  
  
 LG CTAGCCACATCTAGATGCATGTACATAAATTATTGCATACCAAACGATTAAATTAAGCATAAAAAGGAAAAAAA 1472  
 WT CTAGCCACATCTAGATGCATGTACATAAATTATTGCATACCAAACGATTAAATTAAGCATAAAAAGGAAAAAAA  
  
 LG TACTCACATATCTCGACGTAAGATCAATGATATAGTATTTAGATATGCAATATTTATCTTACATCTAAACCTTCTTTCAT 1552  
 WT TACTCACATATCTCGACGTAAGATCAATGATATAGTATTTAGATATGCAATATTTATCTTACATCTAAACCTTCTTTCAT  
  
 LG TCTTAAATATAAGACATTTGTAAGATTTCACTATGGACAACATACGAAACAAAAATCAGTGGATCTCTCTATGCATTTCATT 1632  
 WT TCTTAAATATAAGACATTTGTAAGATTTCACTATGGACAACATACGAAACAAAAATCAGTGGATCTCTCTATGCATTTCATT  
  
 LG ATGTAGTCTATAATAAAATCTTTAAAAAGATCGTATATTTTGCAACGGAGGGAGTAAAAACATAAATTTTAAATAGTAATGT 1712  
 WT ATGTAGTCTATAATAAAATCTTTAAAAAGATCGTATATTTTGCAACGGAGGGAGTAAAAACATAAATTTTAAATAGTAATGT

## FIGURE 12D

LG TGCACGGCTCCACACTCGCAGACGTACCTGCCGAGCCAGATGCCGGGGCGCTGAAGCCGTACCGCGACGACGAGCTCCG 1792  
 WT TGCACGGCTCCACACTCGCAGACGTACCTGCCGAGCCAGATGCCGGGGCGCTGAAGCCGTACCGCGACGACGAGCTCCG  
  
 LG GAAACCTGCTGGCGACGACCGAGAGGGCCCGTACCAGAGCACGACCGCATCTACCGCTACGACGTCTACAAACGACCTCG 1872  
 WT GAAACCTGCTGGCGACGACCGAGAGGGCCCGTACCAGAGCACGACCGCATCTACCGCTACGACGTCTACAAACGACCTCG  
  
 LG GCGAGGGCCGCCCATCCTCGGGGGAACCTCCGACCAACCCTTACCCGCGCGCGCGCGCACGGAGCGCAAGCCCAACGCC 1952  
 WT GCGAGGGCCGCCCATCCTCGGGGGAACCTCCGACCAACCCTTACCCGCGCGCGCGCGCACGGAGCGCAAGCCCAACGCC  
  
 LG AGCGACCCGAGCCTGGAGAGCCGGCTGTGCTGTGGAGCAGATCTACGTGCCGCGGGAAGAGATTTCGGCCACCTCAA 2032  
 WT AGCGACCCGAGCCTGGAGAGCCGGCTGTGCTGTGGAGCAGATCTACGTGCCGCGGGAAGAGATTTCGGCCACCTCAA  
  
 LG GACGTCCGACTTCCTGGGCTACTCCATCAAAGGCCATCACGCAGGGCATCCTGCCGGCCGTGCGCACCTACGTGGACACCA 2112  
 WT GACGTCCGACTTCCTGGGCTACTCCATCAAAGGCCATCACGCAGGGCATCCTGCCGGCCGTGCGCACCTACGTGGACACCA  
  
 LG CCCCCGGGAGTTCGACTCCTTCCAGGACATCATCAACCTCTATAGGGGGGCATCAAGCTGCCCAAGGTGGCCGCCCTG 2192  
 WT CCCCCGGGAGTTCGACTCCTTCCAGGACATCATCAACCTCTATAGGGGGGCATCAAGCTGCCCAAGGTGGCCGCCCTG  
  
 LG GAGGAGTCCCGTAAGCAGTTCCTCCGCTCCAGCTCATCAAGGACCTCCTCCCGCTCGGCGGCGACTCCCTGCTTAAGTCCC 2272  
 WT GAGGAGTCCCGTAAGCAGTTCCTCCGCTCCAGCTCATCAAGGACCTCCTCCCGCTCGGCGGCGACTCCCTGCTTAAGTCCC  
  
 LG CGTGCCCCACATCATCCAGGAGAACAGCAGGCGTGGAGGACCGACGAGGAGTTCCGACGGGAGGTGCTCGCCCGACGTCA 2352  
 WT CGTGCCCCACATCATCCAGGAGAACAGCAGGCGTGGAGGACCGACGAGGAGTTCCGACGGGAGGTGCTCGCCGGCGTCA

(Mutant site) \*

## FIGURE 12E

LG ACCCGGTGATGATCAGCGGTCTCAGCGTGAGTCAGCGATTATTTGTTTCATTGTGTGTATGGTGTCCATGGTGAGAAAG 2432  
 WT ACCCGGTGATGATCAGCGGTCTCAGCGTGAGTCAGCGATTATTTGTTTCATTGTGTGTATGGTGTCCATGGTGAGAAAG  
  
 LG TGCAGATCTTGATTGCGTTGGTTCGATGCACGCGATGCTGCATGCATGCAGGAGTTCCCGCCAAAAGTAGTCTGGACC 2512  
 WT TGCAGATCTTGATTGCGTTGGTTCGATGCACGCGATGCTGCATGCATGCAGGAGTTCCCGCCAAAAGTAGTCTGGACC  
  
 LG CTAGCAAGTTTGGTGACCCACACAGACCACTCATCAGCGGGAGCACATAGAGAAAGAACCTCGAGGGCCTCACGGTGCAGCAG 2592  
 WT CTAGCAAGTTTGGTGACCCACACAGACCACTCATCAGCGGGAGCACATAGAGAAAGAACCTCGAGGGCCTCACGGTGCAGCAG  
  
 LG GTAAATTGGTCCAGCCATCGACATCAACTATGATTTACCTAGGAGTAATTGGTAGCTGTAGATAAATTTGGCTTCGTTGCA 2672  
 WT GTAAATTGGTCCAGCCATCGACATCAACTATGATTTACCTAGGAGTAATTGGTAGCTGTAGATAAATTTGGCTTCGTTGCA  
  
 LG ATTAATTGATGCTGGCCGATCAAGTGATCGTATTGGGTTTGAAATTTGCAGGGCGCTGGAAAGCAACAGGCTGTACATCC 2752  
 WT ATTAATTGATGCTGGCCGATCAAGTGATCGTATTGGGTTTGAAATTTGCAGGGCGCTGGAAAGCAACAGGCTGTACATCC  
  
 LG TTGATCACCATGACCGGTTTCATGCCGTTCCGTGATCGACGTCAACACACCTGCCCGGCAACTTCATCTACGCCACGAGGACC 2832  
 WT TTGATCACCATGACCGGTTTCATGCCGTTCCGTGATCGACGTCAACACACCTGCCCGGCAACTTCATCTACGCCACGAGGACC  
  
 LG CTCTTCTTCTGCGCGCGACGGCAGGCTCAGCGCGCTCGCCATCGAGCTGAGCGAGCCCATCATCCAGGGCGGCTTAC 2912  
 WT CTCTTCTTCTGCGCGCGACGGCAGGCTCAGCGCGCTCGCCATCGAGCTGAGCGAGCCCATCATCCAGGGCGGCTTAC  
  
 LG CACGCCAAGAGCAAGGTTTACACGCCGGTGCCCGAGCGGCTCCGTCTGAAGGCTGGGTGTGGGAGCTCGCCAAAGGCTTACG 2992  
 WT CACGCCAAGAGCAAGGTTTACACGCCGGTGCCCGAGCGGCTCCGTCTGAAGGCTGGGTGTGGGAGCTCGCCAAAGGCTTACG  
  
 LG TCGCCGTCAATGACTCCGGGTGGCACCAGCTCGTCAGCCACTGGTACGTTCTCCACGGTCGATGTGATTTCAGTCAGTCGA 3072  
 WT TCGCCGTCAATGACTCCGGGTGGCACCAGCTCGTCAGCCACTGGTACGTTCTCCACGGTCGATGTGATTTCAGTCAGTCGA



## FIGURE 12F

**LG** TGCACAACTGATCGAAATATGATTGATTGAAACGCCAGGCTGAACACTCACGCGGTGATGGAGCCGTTCTGTGATCT 3152  
**WT** TGCACAACTGATCGAAATATGATTGATTGAAACGCCAGGCTGAACACTCACGCGGTGATGGAGCCGTTCTGTGATCT

**LG** CGACGAACCGGCACCTTAGCGTGACGCACCCGGTGCAAGCTGCTGAGCCCGCACTAC<sup>4</sup>CCGCGACACCATGACCATCAAC 3232  
**WT** CGACGAACCGGCACCTTAGCGTGACGCACCCGGTGCAAGCTGCTGAGCCCGCACTACCCGCGACACCATGACCATCAAC

**LG** GCGCTGGCGCGGAGACGCTCATCAACGCCCGGGCGCATCTTCGAGATGACGGTGTTC<sup>4</sup>CCCGGGCAAGTTCGCGTTGGGGAT 3312  
**WT** GCGCTGGCGCGGAGACGCTCATCAACGCCCGGGCGCATCTTCGAGATGACGGTGTTC<sup>4</sup>CCCGGGCAAGTTCGCGTTGGGGAT

**LG** GTCGGCCGTGGTGTAAGGACTGGAAGTTCA<sup>4</sup>CCGAGCAGGACTGCCGGACGATCTCATCAAGAGGTACGTACCTGGTA 3392  
**WT** GTCGGCCGTGGTGTAAGGACTGGAAGTTCA<sup>4</sup>CCGAGCAGGACTGCCGGACGATCTCATCAAGAGGTACGTACCTGGTA

**LG** AATGTTATGAATGTGTA<sup>4</sup>AAACAAATTTGGCGTCTCGCTCACTGACAGGAACGTGGTAA<sup>4</sup>AAAAAATGCAGGGGCAATGGCGG 3472  
**WT** AATGTTATGAATGTGTA<sup>4</sup>AAACAAATTTGGCGTCTCGCTCACTGACAGGAACGTGGTAA<sup>4</sup>AAAAAATGCAGGGGCAATGGCGG

**LG** TGGAGGACCCGTCGAGCCCGTACAAGGTGCGGTTGCTGGTGTCTGGACTACCCGTACGCGCGGACGCGGCTGGCGATCTGG 3552  
**WT** TGGAGGACCCGTCGAGCCCGTACAAGGTGCGGTTGCTGGTGTCTGGACTACCCGTACGCGCGGACGCGGCTGGCGATCTGG

**LG** CACGCCATTGAGCAGTACGTGAGCGAGTACCTGGCCATCTACTACCCGAACGACGGCGTGTGCAGGGCGGATACGGAGGT 3632  
**WT** CACGCCATTGAGCAGTACGTGAGCGAGTACCTGGCCATCTACTACCCGAACGACGGCGTGTGCAGGGCGGATACGGAGGT


**LG** GCAGGCGTGGTGAAGGAGACGCGCGAGGTCTGGGCACGGCGACCTCAAGGACGCCCCCATGGTGGCCCCAAGATGCAAAAGTG 3712  
**WT** GCAGGCGTGGTGAAGGAGACGCGCGAGGTCTGGGCACGGCGACCTCAAGGACGCCCCCATGGTGGCCCCAAGATGCAAAAGTG

## FIGURE 12G

LG TGCCGGAGCTGGCCAAAGGCGTGACCAACCATCATCTGGATCGGGTCGGCGCTGCATGCGGCAGTCAACTTCGGGCAGTAC 3792  
 WT TGCCGGAGCTGGCCAAAGGCGTGACCAACCATCATCTGGATCGGGTCGGCGCTGCATGCGGCAGTCAACTTCGGGCAGTAC  
  
 LG CCCTACGCGGGGTTCTCTCCGAACCGGCCGACGGTGAGCCGGCCCGCATGCCGAGCCCCGGCACGGAGGAGTACGCGGA 3872  
 WT CCCTACGCGGGGTTCTCTCCGAACCGGCCGACGGTGAGCCGGCCCGCATGCCGGAGCCCCGGCACGGAGGAGTACGCGGA  
  
 LG GCTGGAGCGGACCCGGAGCGGGCCCTTCATCCACACCATCACGAGCCAGATCCAGACCATCATCGGCGTGTGCTGTGG 3952  
 WT GCTGGAGCGGACCCGGAGCGGGCCCTTCATCCACACCATCACGAGCCAGATCCAGACCATCATCGGCGTGTGCTGTGG  
  
 LG AGGTGCTGTGAAAGCACTCTCCGACGAGCTGTACCTCGGGCAGCGGGACACGCCGGAGTGGACCTCGGACCCAAAGGCC 4032  
 WT AGGTGCTGTGAAAGCACTCTCCGACGAGCTGTACCTCGGGCAGCGGGACACGCCGGAGTGGACCTCGGACCCAAAGGCC  
  
 LG CTGGAGGTGTTCAAGCGGTTACGCGACCGGCTGGTGGAGATCGAGAGCAAGGTGGTGGCATGAACCATGACCCGGAGCT 4112  
 WT CTGGAGGTGTTCAAGCGGTTACGCGACCGGCTGGTGGAGATCGAGAGCAAGGTGGTGGCATGAACCATGACCCGGAGCT  
  
 LG CAAGAACCGCAACGGCCCGGCTAAGTTTCCCTACATGCTGCTCTACCCCAACACCTCCGACCCACAAGGGCGCGTGCCG 4192  
 WT CAAGAACCGCAACGGCCCGGCTAAGTTTCCCTACATGCTGCTCTACCCCAACACCTCCGACCCACAAGGGCGCGTGCCG  
  
 translation stop (4229)  
 LG GGCTTACCGCCAAAGGCGATCCCCAACAGCATCTCCATCTAATCTAAGCCATCGGCAACCATGGATGAATAAAGGGCGTTC 4272  
 WT GGCTTACCGCCAAAGGCGATCCCCAACAGCATCTCCATCTAATCTAAGCCATCGGCAACCATGGATGAATAAAGGGCGTTC  
  
 LG GCCACGTACGAAACTTGTGAGAGAGATTGGTGTAGTGTGTCTGTGACAGTACTATGTACGAGTTGCTCTTTAAGCCGA 4352  
 WT GCCACGTACGAAACTTGTGAGAGAGATTGGTGTAGTGTGTCTGTGACAGTACTATGTACGAGTTGCTCTTTAAGCCGA  
  
 LG ATAAATAAAGCAGATTTGCTTCC 4375  
 WT ATAAATAAAGCAGATTTGCTTCC

FIGURE 13

Wild-type *LOX-1* gene structure:



Line G *lox-1* gene structure:





FIGURE 14




**FIGURE 15**

FIGURE 16




FIGURE 17






FIGURE 18





**FIGURE 21**





FIGURE 20

Backcrossing program for Line G to cv Alexis



**FIGURE 22 A**  
**LIPOXYGENASE AMINO ACID SEQUENCE ALIGNMENT**

```

1          E1      H          60
Gm1 MFS.....A.GHKIKGTVVLMRKNELE.....VNPDGSAV
Gm2 MFSVPGVSGIILNRGG.GHKIKGTVVLMRKNVLDNFNSVADLTGKNVGGGLIGTGLNVVGSTL
Gm3 M.....LGGLLHRG...HKIKGTVVLMRKNVLDVNSVT....SVGGIIGQGLDLVGSTL
Hv2 MLGVGGIVSDLTGGIRGAHLKGSVLMRKNALDFN.....DFGAHVM
Hv1 ML.LGGLIDTLTGANKSARLKGTVLMRKNVLDLN.....DFGATII
    61      +      E2      ++ +      E3      +      +      E4 120
Gm1 DNLNAFLGRSVSLQLISATKADAH..GKGKVGKDTFLEG..INTSLPTLGAGESAFNIHF
Gm2 DNLTAFLGRSVALQLISATKPLAN..GKGKVGKDTFLEG..IIVSLPTLGAGESAFNIQF
Gm3 DTLTAFLGRSVSLQLISATKADAN..GKGKLGKATFLEG..IITSLPTLGAGQSAFKINF
Hv2 DGVTELLGRGVTCQLISSTNVDHNNNGRGKVGAEANLEQWLLPTNLPFITTGENKFAVTF
Hv1 DGIGEF LGKGVTCQLISSTAVDQDNGGRGKVGAEAELEQWV..TSLPSLTGSGKFGLTF
    121      + +      E5      E6      +      E7      E8 180
Gm1 EWD.GSMGIPGAFYIKNYMQVEFFLKSLTLEAISNQ.GTIRFVCNSWVYNTKLYKSVRIF
Gm2 EWD.ESMGIPGAFYIKNYMQVEFFLKSLTLEDVNPQ.GTIRFVCNSWVYNTKLYKSVRIF
Gm3 EWD.DSGSIPGAFYIKNFMQTEFFLVSLTLEDIPNH.GSIHFVCNSWIYNAKLFKSDRIF
Hv2 DWSVDKLGVPGAIIIVKNNHASEFFLKTITLDNVPGR.GTIVFVANSWVYPQAKYRYNRVF
Hv1 DWEVEKLGVPGAIVVNNYHSSEFFLKTITLHDVPGRSGNLTFFVANSWIYPAANYRYSRVF
    181      +      +      H1      +      +      E      +      +      240
Gm1 FANHTYVPSETPAPLVSYREEELKSLRGNGT.GERKEYDRIYDYDVYNDLGNPDKSEKLA
Gm2 FANHTYVPSETPAALVGYREEELKNLRGDGK.GERKEHDRIYDYDVYNDLGNPDHGENFA
Gm3 FANQTYLPSETPAPLVKYREEELHNLRGDGT.GERKEWERIYDYDVYNDLGD PDKGENHA
Hv2 FANDTYLPHQMPAALKPYRDELRNLRGDDQQGPYLDHDRVYRYDVYNDLGD.....S
Hv1 FANDTYLPSQMPAALKPYRDELRNLRGDDQQGPYQEHDRYRYDVYNDLGE.....G
    241 ++      +      +      +      •••      • 300
Gm1 RPVLGGSSTFFVPRRGRTGRGPTVTDPNTEKQ....GEVFYVPRDENLGHLSKDAFEIG
Gm2 RPILGGSSTHEVPRRGRTGRYPTRKQNSEKP....GEV.YVPRDENFGHLKSSDFEAYG
Gm3 RPVLGGNDTFVPRRGRTGRKPTRKDPNSESER....SNDVYLPRDEAFGHLKSSDFETYG
Hv2 RDVLGGSKDLFVPRRCRTGRKPSDSKPDHESRLLLLVQNVYVLRDELFGHLKQSDLLGYT
Hv1 RPILGGSNDHFPVPRGRTERKPNASDPSLESRLSLL.EQIYVPRDEKFGHLKTSDFEIGYS
    •      H2      +      H3      ++      +      H4      +      360
Gm1 TKLSLQIVQPAFESAFDLKSTPIEFHFSQDQVHDLYEGGIKLP....DVISTIIPLPVIK
Gm2 IKSLSQYVLPAFESVFDLNFPTNEFDSFQDVRDLHEGGIKLPT....EVISTIMPLPVVK
Gm3 LKSVSQNVLPQLQSAFDLNFPTREFDSFDEVHGLYSGGIKLP....DIISKISPLPVLK
Hv2 LKGWLDGIILAIRTYVDL..SPGEFDSFADILKLYEGGIKLPNIPALEEVRKRFPQLQVK
Hv1 IKAITQGILPAVRTYVDT..TPGEFDSFQDIINLYEGGIKLPKVALEELRKQFPLQLIK
    E10 +E11 + +      H6      ♦      +      E12      ++      420
Gm1 ELYRTDGHILKFPQPHVVQVSQSAAWMTDEEFAREMIAGVNPCVIRGLEEFPPKSNLDPA
Gm2 ELFRTDGEQVLKFPFPPHVIQVSKSAWMTDEEFAREMIVAGVNPVIRGLQEFPPKSNLDPT
Gm3 EIFRTDGEQALKFPFPPKVIQVSKSAWMTDEEFAREMLAGVNPVIRCLKDFPPRSKLDSDQ
Hv2 DLIPKGGDFLLKLPKPEIIKVQKAWMTDEEFAREMLAGVNPVMIKRLTEFPKSTLDPS
Hv1 DLLPVGDSLLKLPVPHIIQENKQAWMTDEEFAREVLAGVNPVMITRLTEFPKSSLDPS
    +      +      +      H7      E13      H8      480
Gm1 IYGDQSSKITADSLD..LDGYTMDEALGSRRLFMDYHDIEMPYVRQINQLNSAKTYATR
Gm2 IYGEQTSKITADALD..LDGYTVDEALASRRLFMDYHDVEMPYIRRINQ.TYAKAYATR
Gm3 VYGDHTSQITKEHLEPNLEGLTVDEAIQNKRLFMDHHDPIIMPYLRRINA.TSTKAYATR
Hv2 KYGDHTSTMTEEHVAKSLEGLTVQQAALAGNRLYIVDQHDNLMFLIDINNLDASFYATR
Hv1 KFGDHTSTITAEHIEKNLEGLTVQQALESNRLYIHDHDFMPFLIDVNNLPGNFIYATR

```

FIGURE 22 B

	E14	+	E15	+	E16	+	+	H9	540
Gm1	TILFLREDGTLKPVAIELSLPHSAGDLSAAVSQVVLPAKEG.VESTIWLILAKAYVIVNDS								
Gm2	TILFLRENGTLKPVAIELSLPHAGDLSGAVSQVILPAKEG.VESTIWLILAKAYVVVNDG								
Gm3	TILFLKNDGTLRPLAIELSLPHPGDQSGAFSQVFLPADEG.VESSIWLILAKAYVVVNDG								
Hv2	TLLFLRGDGTLPVAIELSSPLIQGELTTAKSAVYTPQHAG.VEGWIWQLAKAYASVNDY								
Hv1	TLFFLRGDGRLTPLAIELSEPIIQGGLTTAKSKVYTPVPSGSVEGWVWELAKAYVAVNDS								
	541	•	•	•	H9		+H10	+	•
									H11
									600+
Gm1	CYHQLMSHWLNTHAAMEPFVIATHRHLSVLHPYKLLTPHYRNNMIVNALAROSLINANG								
Gm2	CYHQLMSHWLNTHAVIEPFIATNRHLSALHPYKLLTPHYRDTMIVNALAROSLINADG								
Gm3	CYHQLVSHWLNTHAVVEPFIATNRHLSVVHPYKLLHPHYRDTMIVNGLARLSLVNDGG								
Hv2	GWHLQISHWLNTHAVMEPFVIATNRQLSVTHPVYKLLHPHYRDTMIVNARARGLLINAGG								
Hv1	GWHLQVSHWLNTHAVMEPFVISTNRHLSVTHPVHKLLSPHYRDTMTIVNALARQTLINAGG								
	601	H12		H13		+H14	+	E	+
								E	+660
Gm1	IIETTFELPSKYSVEMSSAVYKNWVFTDQALPADLIKRGVAIKDPSTPHGVRLIEDYPYA								
Gm2	IIEKSEFLPSKHSVEMSSAVYKNWVFTDQALPADLIKRGVAIKDPSAPHGLRLIEDYPYA								
Gm3	VIEQTFELWGRYSVEMSSAVYKDWVFTDQALPADLIKRGMAIEDPSCPHGIRLVIEDYPYT								
Hv2	VIEMTVFPKHAMPMSMVYKHWNFTEQALPADLIKRGMAVEDASSPHKVRLLIKDYPIYA								
Hv1	IFEMTVFPGKFALGMSAVYKDWKFTEQGLPDDLIKRGMAVEDPSSPYKVRLLVSDYPYA								
	661	H15		H16		H17	+	+	+
									720
Gm1	ADGLEIWAIAIKTWVQEVVPLYARDDVDKNDSELQHWKEAVEKGHGDLKDKPWWPKLQT								
Gm2	VDGLEIWAIAIKTWVQEVVSLYARDDVDKPDSELQQWKEAVEKGHGDLKDKPWWPKLQT								
Gm3	VDGLEIWDIAIKTWVHEYVFLYKSDDTLREDPELQACWKEAVEVGHGDKKNEPWWPKMQT								
Hv2	TDGLAVWDIAEQWVSDYLTIIYPNDGVLQGDVELQAWWKEVREVGHGDLKDAAWPKMQT								
Hv1	ADGLAIWHAIEQYVSEYLAIIYPNDGVLQGDTEVQAWWKETREVGHGDLKDAFWPKMQS								
	721		H18	•	•	•	+	•	+
									H19
									780
Gm1	LEDLVECLIIIIWTASALHAAVNFGQYPYGLIIMNRPTASRRLLPEKGTPEYEEMINNHE								
Gm2	IEELVEICTIIIIWTASALHAAVNFGQYPYGFIIINRPTSSRRLLPEKGTPEYEMVKS HQ								
Gm3	REELVEACIIIIWTASALHAAVNFGQYPYGLIINRPTLSRRFMPEKGS AEYELRKNPQ								
Hv2	VAELIKACATIIWTGSALHAAVNFGQYPYSGYHPNKPESASRRPMPVQGS EYAE LERDPE								
Hv1	VP ELAKACTIIWIGSALHAAVNFGQYPYAGFLPNRPTVSRRMPPEPGTEEYAE LERDPE								
	781	H20		H21	•	•	•	+	+
									H22
									840
Gm1	KAYLRTITSKLPTLISLSVIEIILSTHASDEVYLGQRDNPHWTS DSKALQAFQKFGNKLKE								
Gm2	KAYLRTITSKFQTLVDLSVIEIILSRHASDEVYLGQRDNPHWTS DSKALQAFQKFGNKLKE								
Gm3	KAYLKTITPKFQTLIDLSVIEIILSRHASDEVYLGQRDNPNWTS DTRALEAFKRFGNKLAQ								
Hv2	KAFIRTITSQFHALVGISLMEIILSKHSSDEVYLGQHDTPAWTS DAKALEAFKRFGAKLEG								
Hv1	RAFIHTITSQIQTIIIGVSLLEVILSKHSSDELYLGQRDTPWTS DPKALEVFKRFS DRLVE								
	841		++	+	+	+	+	+	+
									895*
Gm1	IEEKLVRNNDPSLQGNRLGPVQLPYTLLYPSS EEE.....GLTFRGIPNSIS I								
Gm2	IEEKLARKNNDQSL.SNRLGPVQLPYTLLHPNSE.....GLTCRGIPNSIS I								
Gm3	IENKLSERNNDEKLR.NRCGPVQMPYTLLLPSSKE.....GLTFRGIPNSIS I								
Hv2	IEKQVAMNSDPQLK.NRTGPAKFPPYMLLYPNTSDHTGQAEGLTARGIPNSIS I								
Hv1	IESKVVGMMHDP ELK.NRNGPAKFPPYMLLYPNTSDHKGAAAGLTAKGIPNSIS I								

Gm1 SEQ ID NO:15

Gm2 SEQ ID NO:16

Gm3 SEQ ID NO:17

Hv2 SEQ ID NO:18

Hv1 SEQ ID NO:9

•The numbering system shown above the sequence alignment does not correspond to the actual amino acid number of any lipoxxygenase sequence in the alignment.

SEQLIST.ST25  
SEQUENCE LISTING

<110> Douma, Anneke  
Doderer, Albert  
Cameron-Mills, Verena  
Skadhauge, Birgitte  
Bech, Lene

<120> LOW LIPOXYGENASE 1 BARLEY

<130> 11225.11W001

<160> 18

<170> PatentIn version 3.0

<210> 1

<211> 143

<212> DNA

<213> Hordeum vulgare

<400> 1  
ctaagccatc ggcaaccatg gatgaataaa gggcgttcgc cacgtacgaa acttgctcgag 60  
agattggtgt agtgtgtgtc tgtgacagta ctatgtcagc agttgctctt taagccgaat 120  
aaataaagca gatttgcttc caa 143

<210> 2

<211> 24

<212> DNA

<213> Hordeum vulgare

<400> 2  
gaaaagcttg gaggtagacg ctgc 24

## SEQLIST.ST25

<210> 3  
<211> 33  
<212> DNA  
<213> Hordeum vulgare

<400> 3  
tataggatcc ttgttcttgg cctcctctcc tcg 33

<210> 4  
<211> 22  
<212> DNA  
<213> Hordeum vulgare

<400> 4  
agtgaaaaac agtgtgctgg tg 22

<210> 5  
<211> 21  
<212> DNA  
<213> Hordeum vulgare

<400> 5  
ggcttaaaga gcaactgctg a 21

<210> 6  
<211> 23  
<212> DNA  
<213> Hordeum vulgare

<400> 6  
caagatgcat atgctgctgg gag 23

<210> 7  
<211> 29  
<212> DNA

## SEQLIST.ST25

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 7

cgatggttta aattagatgg agatgctgt

29

&lt;210&gt; 8

&lt;211&gt; 4663

&lt;212&gt; DNA

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 8

cagccccatg catgcacatg cacatgcaca tgcacatgca gtgcagccaa gcaccgctcg	60
atgggcgatc acccgtcacg ggaccggagc gcgccatgcg aagcacgagg agggcacgtc	120
accgtccgcg cgcagcacgt ggagagcacg tcgccgtccg atccatctct ccaaagccga	180
gcgccacacc accgggaccg gaccgggacc ggcctataaa ttgcccggac cgagctgcaa	240
gcagctcctc acacacactc acgcaacaca catccatctt cactgaaaag tgaaaaacag	300
tgtgctggtg ccattggttg gagcagtga aagcaggaga ggaggccaag aacaagatgc	360
tgctgggagg gctgatcgac accctcacgg gggcgaacaa gagcgcccg ctcaaggga	420
cggtggtgct catgcgcaag aacgtgctgg acctcaacga cttcggcgcc accatcatcg	480
acggcatcgg cgagttcctc ggcaagggcg tcacctgcca gcttatcagc tccaccgccg	540
tcgaccaagg taatcactac cctcctccgg ctttctcctc tgtttacaag atatagtatt	600
tctttcgtgt gggccggcgg ccatggatgg atggatgtgt ctggatcggc taaagaagat	660
aggatagcta gccctggccg gtcgtcttta cctgagcatg ggcataatgcc atcgaaaaaa	720
gagacaacag catgcatgca tgggtgcgcg accagaccac gcagagcacc ggatgctcga	780
gacaaagcaa cacaacaagc aaggacgaca cgtcaaaagc aacacaacaa gcaaggacgg	840
cacgtcaaaa gcaacacaaa cctaaactaa agcacaaaga cgtaagagca agcacacaat	900
cagcaggcta taaacagttg tcatcaaaaa caacgctgga agagagagag aaggaaggaa	960
gtagtagcca tgaaaaatta aatcaccggg cgttgctctt tgcccaacaa ttaatcaagc	1020
aggggtacgtg gcatgtatag ttcttgtaag taaactaagc atgtgatatg agaaggtacg	1080
tgggtggtgca gacaacggcg gtcgcgggaa ggtgggagcg gaggcggagc tggagcagtg	1140
ggtgacgagc ctgccgtcgc tgacgacggg ggagtccaag ttcggcctca ccttcgactg	1200
ggaggtggag aagctcgggg tgccgggagc catcgtcgtc aacaactacc acagctccga	1260
gttcctgctt aaaaccatca ccctccacga cgtccccggc cgcagcggca acctcacctt	1320



## SEQLIST.ST25

cgtcgccaac	tcatggatct	accccgccgc	caactaccga	tacagccgcg	tcttcttcgc	1380
caacgacgtg	cgtggatttt	cctctacttt	cctctccttt	cattttcacc	gccttcgtca	1440
ttcatggtcg	atcattaagt	cttgccagga	caatagatga	tgagctagga	gtggttacca	1500
cttagcagta	cgtacattat	ttattccgtg	ttggtagaaa	aggatatggt	ttggtgcaga	1560
tcgacacaag	attgaatgaa	agttgcaccg	tggcaccgtg	gcagcgtggt	aggtgaaaat	1620
aactgttgca	cggatccacc	cacatgattg	ttttcatgaa	taaacttttt	aaggatgtgt	1680
ctagccacat	ctagatgcat	gtcacataat	tattgcatac	caaaacgatt	aaattaagca	1740
taaaaagaaa	aggaaaaaaa	tactcacata	tctcgacgta	agatcaatga	tatagtattt	1800
agatatgcaa	tatttatctt	acatctaaac	ctttcttcat	tcttaaatat	aagacatttg	1860
taagatttca	ctatggacaa	catacgaaac	aaaatcagtg	gatctctcta	tgcattcatt	1920
atgtagtcta	taataaaatc	tttaaaagat	cgtatatattt	gcaacggagg	gagtaaaaca	1980
taacttttta	atagtaatgt	tgcacggctc	cacactcgca	gacgtacctg	ccgagccaga	2040
tgccggcggc	gctgaagccg	taccgcgacg	acgagctccg	gaacctgcgt	ggcgacgacc	2100
agcagggccc	gtaccaggag	cacgaccgca	tctaccgcta	cgacgtctac	aacgacctcg	2160
gcgagggccg	ccccatcctc	ggcggaact	ccgaccaccc	ttacccgcgc	cgcgcccgca	2220
cggagcgcaa	gcccaacgcc	agcgacccga	gcctggagag	ccggctgtcg	ctgctggagc	2280
agatctacgt	gccgcgggac	gagaagttcg	gccacctcaa	gacgtccgac	ttcctgggct	2340
actccatcaa	ggccatcacg	cagggcatcc	tgccggccgt	gcgcacctac	gtggacacca	2400
ccccggcgga	gttcgactcc	ttccaggaca	tcatcaacct	ctatgagggc	ggcatcaagc	2460
tgcccaaggt	ggccgccctg	gaggagctcc	gtaagcagtt	cccgtccag	ctcatcaagg	2520
acctcctccc	cgtcggcggc	gactccctgc	ttaagctccc	cgtgccccac	atcatccagg	2580
agaacaagca	ggcgtggagg	accgacgagg	agttcgcacg	ggaggtgctc	gccggcgtca	2640
accgggtcat	gatcacgcgt	ctcacggtga	gtcagcgatt	atttgttcat	tgtgtgtgta	2700
tggtgtccat	ggtgagaaa	tgcatatctt	gatttgcggt	gggtcgcatg	cacgcatgct	2760
gcatgcatgc	aggagtcccc	gccaaaaagt	agtctggacc	ctagcaagtt	tggtgaccac	2820
accagcacca	tcacggcgga	gcacatagag	aagaacctcg	agggcctcac	ggtgcagcag	2880
gtaattgggtc	caagccatcg	acatcaacta	tgatttacct	aggagtaatt	ggtagctgta	2940
gataatttgg	cttcgttgca	attaatttga	tgctggccga	tcaagtgatc	gtattgggtt	3000
tgaaatttgc	aggcgtgga	aagcaacagg	ctgtacatcc	ttgatcacca	tgaccggttc	3060
atgccgttcc	tgatcgacgt	caacaacctg	cccggcaact	tcatctacgc	cacgaggacc	3120
ctcttcttcc	tgcgcggcga	cggcaggctc	acgccgctcg	ccatcgagct	gagcgagccc	3180
atcatccagg	gcggccttac	cacggccaag	agcaaggttt	acacgccggt	gccagcggc	3240

## SEQLIST.ST25

```

tccgtcgaag gctgggtgtg ggagctcgcc aaggcctacg tcgccgtcaa tgactccggg 3300
tggcaccagc tcgtcagcca ctggtacgtt ctccacggtc gatgtgattc agtcagtcga 3360
tgcacaacaa ctgatcgaaa tatgattgat tgaaacgcgc aggctgaaca ctcacgcggg 3420
gatggagccg ttcgtgatct cgacgaaccg gcaccttagc gtgacgcacc cgggtgcacaa 3480
gctgctgagc ccgcactacc gcgacaccat gaccatcaac gcgctggcgc ggcagacgct 3540
catcaacgcc ggcgcatct tcgagatgac ggtgttcccc ggcaagtctg cgttggggat 3600
gtcggccgtg gtgtacaagg actggaagtt caccgagcag ggactgccgg acgatctcat 3660
caagaggtac gtacctggtg aatgttatga atgtgtaaaa caaattgggc gtctcgctca 3720
ctgacaggaa cgtggtaaaa aaaatgcagg ggcattggcg tggaggaccc gtcgagccc 3780
tacaaggtgc ggttgctggt gtcggactac ccgtacgcgc cggacgggct ggcgatctgg 3840
cacgccattg agcagtacgt gagcgagtac ctggccatct actacccgaa cgacggcgtg 3900
ctgcagggcg atacggaggt gcaggcgtgg tggaaggaga cgcgcgaggt cgggcacggc 3960
gacctcaagg acgccccatg gtggcccaag atgcaaagtg tgccggagct ggccaaggcg 4020
tgcaccacca tcatctggat cgggtcggcg ctgcatgcgc cagtcaactt cgggcagtac 4080
ccctacgcgc ggttcctccc gaaccggccg acggtgagcc ggcgccgcac gccggagccc 4140
ggcacggagg agtacgcgga gctggagcgc gacccggagc gggccttcat ccacaccatc 4200
acgagccaga tccagacat catcggcgtg tcgctgctgg aggtgctgtc gaagcactcc 4260
tccgacgagc tgtacctcg gcagcgggac acgccggagt ggacctcgga cccaaaggcc 4320
ctggaggtgt tcaagcggtt cagcgaccgc ctggtggaga tcgagagcaa ggtggtgggc 4380
atgaaccatg acccgagct caagaaccgc aacggcccgc ctaagtctcc ctacatgctg 4440
ctctacccca acacctccga ccacaaggcg gccgtgccg ggcttaccgc caagggcatc 4500
cccaacagca tctcatcta atctaagcca tcggcaacca tggatgaata aagggcggtc 4560
gccacgtacg aaacttgctg agagattggt gtagtggtg tctgtgacag tactatgtca 4620
gcagttgctc ttttaagccga ataaataaag cagatttgct tcc 4663

```

&lt;210&gt; 9

&lt;211&gt; 862

&lt;212&gt; PRT

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 9

Met Leu Leu Gly Gly Leu Ile Asp Thr Leu Thr Gly Ala Asn Lys Ser

SEQLIST.ST25

1	5	10	15
Ala Arg Leu Lys Gly Thr Val Val Leu Met Arg Lys Asn Val Leu Asp	20	25	30
Leu Asn Asp Phe Gly Ala Thr Ile Ile Asp Gly Ile Gly Glu Phe Leu	35	40	45
Gly Lys Gly Val Thr Cys Gln Leu Ile Ser Ser Thr Ala Val Asp Gln	50	55	60
Asp Asn Gly Gly Arg Gly Lys Val Gly Ala Glu Ala Glu Leu Glu Gln	65	70	75
Trp Val Thr Ser Leu Pro Ser Leu Thr Thr Gly Glu Ser Lys Phe Gly	85	90	95
Leu Thr Phe Asp Trp Glu Val Glu Lys Leu Gly Val Pro Gly Ala Ile	100	105	110
Val Val Asn Asn Tyr His Ser Ser Glu Phe Leu Leu Lys Thr Ile Thr	115	120	125
Leu His Asp Val Pro Gly Arg Ser Gly Asn Leu Thr Phe Val Ala Asn	130	135	140
Ser Trp Ile Tyr Pro Ala Ala Asn Tyr Arg Tyr Ser Arg Val Phe Phe	145	150	155
Ala Asn Asp Thr Tyr Leu Pro Ser Gln Met Pro Ala Ala Leu Lys Pro	165	170	175
Tyr Arg Asp Asp Glu Leu Arg Asn Leu Arg Gly Asp Asp Gln Gln Gly	180	185	190
Pro Tyr Gln Glu His Asp Arg Ile Tyr Arg Tyr Asp Val Tyr Asn Asp	195	200	205
Leu Gly Glu Gly Arg Pro Ile Leu Gly Gly Asn Ser Asp His Pro Tyr	210	215	220
Pro Arg Arg Gly Arg Thr Glu Arg Lys Pro Asn Ala Ser Asp Pro Ser	225	230	235
Leu Glu Ser Arg Leu Ser Leu Leu Glu Gln Ile Tyr Val Pro Arg Asp	245	250	255
Glu Lys Phe Gly His Leu Lys Thr Ser Asp Phe Leu Gly Tyr Ser Ile	260	265	270
Lys Ala Ile Thr Gln Gly Ile Leu Pro Ala Val Arg Thr Tyr Val Asp	275	280	285
Thr Thr Pro Gly Glu Phe Asp Ser Phe Gln Asp Ile Ile Asn Leu Tyr	290	295	300
Glu Gly Gly Ile Lys Leu Pro Lys Val Ala Ala Leu Glu Glu Leu Arg	305	310	315
Lys Gln Phe Pro Leu Gln Leu Ile Lys Asp Leu Leu Pro Val Gly Gly	325	330	335
Asp Ser Leu Leu Lys Leu Pro Val Pro His Ile Ile Gln Glu Asn Lys			

**SEQLIST.ST25**

340										345				350			
Gln	Ala	Trp	Arg	Thr	Asp	Glu	Glu	Phe	Ala	Arg	Glu	Val	Leu	Ala	Gly		
		355					360					365					
Val	Asn	Pro	Val	Met	Ile	Thr	Arg	Leu	Thr	Glu	Phe	Pro	Pro	Lys	Ser		
	370					375					380						
Ser	Leu	Asp	Pro	Ser	Lys	Phe	Gly	Asp	His	Thr	Ser	Thr	Ile	Thr	Ala		
385					390					395					400		
Glu	His	Ile	Glu	Lys	Asn	Leu	Glu	Gly	Leu	Thr	Val	Gln	Gln	Ala	Leu		
				405					410					415			
Glu	Ser	Asn	Arg	Leu	Tyr	Ile	Leu	Asp	His	His	Asp	Arg	Phe	Met	Pro		
			420					425					430				
Phe	Leu	Ile	Asp	Val	Asn	Asn	Leu	Pro	Gly	Asn	Phe	Ile	Tyr	Ala	Thr		
		435					440					445					
Arg	Thr	Leu	Phe	Phe	Leu	Arg	Gly	Asp	Gly	Arg	Leu	Thr	Pro	Leu	Ala		
	450					455					460						
Ile	Glu	Leu	Ser	Glu	Pro	Ile	Ile	Gln	Gly	Gly	Leu	Thr	Thr	Ala	Lys		
465					470					475					480		
Ser	Lys	Val	Tyr	Thr	Pro	Val	Pro	Ser	Gly	Ser	Val	Glu	Gly	Trp	Val		
				485					490					495			
Trp	Glu	Leu	Ala	Lys	Ala	Tyr	Val	Ala	Val	Asn	Asp	Ser	Gly	Trp	His		
			500					505					510				
Gln	Leu	Val	Ser	His	Trp	Leu	Asn	Thr	His	Ala	Val	Met	Glu	Pro	Phe		
		515					520					525					
Val	Ile	Ser	Thr	Asn	Arg	His	Leu	Ser	Val	Thr	His	Pro	Val	His	Lys		
	530					535					540						
Leu	Leu	Ser	Pro	His	Tyr	Arg	Asp	Thr	Met	Thr	Ile	Asn	Ala	Leu	Ala		
545					550					555					560		
Arg	Gln	Thr	Leu	Ile	Asn	Ala	Gly	Gly	Ile	Phe	Glu	Met	Thr	Val	Phe		
				565					570					575			
Pro	Gly	Lys	Phe	Ala	Leu	Gly	Met	Ser	Ala	Val	Val	Tyr	Lys	Asp	Trp		
			580					585					590				
Lys	Phe	Thr	Glu	Gln	Gly	Leu	Pro	Asp	Asp	Leu	Ile	Lys	Arg	Gly	Met		
		595					600					605					
Ala	Val	Glu	Asp	Pro	Ser	Ser	Pro	Tyr	Lys	Val	Arg	Leu	Leu	Val	Ser		
	610					615					620						
Asp	Tyr	Pro	Tyr	Ala	Ala	Asp	Gly	Leu	Ala	Ile	Trp	His	Ala	Ile	Glu		
625					630					635					640		
Gln	Tyr	Val	Ser	Glu	Tyr	Leu	Ala	Ile	Tyr	Tyr	Pro	Asn	Asp	Gly	Val		
				645					650					655			
Leu	Gln	Gly	Asp	Thr	Glu	Val	Gln	Ala	Trp	Trp	Lys	Glu	Thr	Arg	Glu		
			660					665					670				
Val	Gly	His	Gly	Asp	Leu	Lys	Asp	Ala	Pro	Trp	Trp	Pro	Lys	Met	Gln		

## SEQLIST.ST25

675                      680                      685  
 Ser Val Pro Glu Leu Ala Lys Ala Cys Thr Thr Ile Ile Trp Ile Gly  
 690                      695                      700  
 Ser Ala Leu His Ala Ala Val Asn Phe Gly Gln Tyr Pro Tyr Ala Gly  
 705                      710                      715                      720  
 Phe Leu Pro Asn Arg Pro Thr Val Ser Arg Arg Arg Met Pro Glu Pro  
 725                      730                      735  
 Gly Thr Glu Glu Tyr Ala Glu Leu Glu Arg Asp Pro Glu Arg Ala Phe  
 740                      745                      750  
 Ile His Thr Ile Thr Ser Gln Ile Gln Thr Ile Ile Gly Val Ser Leu  
 755                      760                      765  
 Leu Glu Val Leu Ser Lys His Ser Ser Asp Glu Leu Tyr Leu Gly Gln  
 770                      775                      780  
 Arg Asp Thr Pro Glu Trp Thr Ser Asp Pro Lys Ala Leu Glu Val Phe  
 785                      790                      795                      800  
 Lys Arg Phe Ser Asp Arg Leu Val Glu Ile Glu Ser Lys Val Val Gly  
 805                      810                      815  
 Met Asn His Asp Pro Glu Leu Lys Asn Arg Asn Gly Pro Ala Lys Phe  
 820                      825                      830  
 Pro Tyr Met Leu Leu Tyr Pro Asn Thr Ser Asp His Lys Gly Ala Ala  
 835                      840                      845  
 Ala Gly Leu Thr Ala Lys Gly Ile Pro Asn Ser Ile Ser Ile  
 850                      855                      860

&lt;210&gt; 10

&lt;211&gt; 2818

&lt;212&gt; DNA

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 10

agtgaaaaac agtgtgctgg tgccattggt tggagcagtg aaagcgagga gaggaggcca 60  
 agaacaagat gctgctggga gggctgatcg acaccctcac gggggcgaac aagagcgccc 120  
 ggctcaaggg cacggtggtg ctcatgcgca agaacgtgct ggacctcaac gacttcggcg 180  
 ccaccatcat cgacggcatc ggcgagttcc tcggcaaggg cgtcacctgc cagcttatca 240  
 gctccaccgc cgtcgaccaa gacaacggcg gtcgcgggaa ggtgggcgcg gaggcggagc 300  
 tggagcagtg ggtgacgagc ctgccgtcgc tgacgacggg ggagtccaag ttcggcctca 360  
 ccttcgactg ggaggtggag aagctcgggg tgccgggcgc catcgtcgtc aacaactacc 420  
 acagctccga gttcctgctt aaaaccatca ccctccacga cgtccccggc cgcagcggca 480  
 acctcacctt cgtcgccaac tcatggatct accccgccgc caactaccga tacagccgcg 540

## SEQLIST.ST25

tcttcttcgc caacgacacg tacctgccga gccagatgcc ggcggcgctg aagccgtacc	600
gcgacgacga gctccggaac ctgctggcg acgaccagca gggcccgtac caggagcacg	660
accgcatcta ccgctacgac gtctacaacg acctcggcga gggccgcccc atcctcggcg	720
gcaactccga ccacccttac ccgcgcgcg gccgcacgga gcgcaagccc aacgccagcg	780
acccgagcct ggagagccgg ctgtcgtgc tggagcagat ctacgtgccg cgggacgaga	840
agttcggcca cctcaagacg tccgacttcc tgggtactc catcaaggcc atcacgcagg	900
gcacctgcc ggccgtgcgc acctacgtgg acaccacccc cggcgagttc gactccttcc	960
aggacatcat caacctctat gagggcggca tcaagctgcc caaggaggcc gccctggagg	1020
agctccgtaa gcagttcccg ctccagctca tcaaggacct cctccccgtc ggcggcgact	1080
ccctgcttaa gctccccgtg cccacatca tccaggagaa caagcaggcg tggaggaccg	1140
acgaggagtt cgcacgggag gtgctcgccg gcgtcaaccc ggtcatgac acgcgtctca	1200
cggagttccc gccaaaaagt agtctggacc ctagcaagtt tggtgaccac accagcacca	1260
tcacggcgga gcacatagag aagaacctcg agggcctcac ggtgcagcag gcgctggaaa	1320
gcaacaggct gtacatcctt gatcaccatg accggttcat gccgttcctg atcgacgtca	1380
acaacctgcc cggcaacttc atctacgcca cgaggacct cttcttcctg cgcggcgacg	1440
gcaggctcac gccgctcgcc atcgagctga gcgagcccat catccagggc ggccttacca	1500
cggccaagag caaggtttac acgccggtgc ccagcggtc cgtcgaaggc tgggtgtggg	1560
agctcgccaa ggcctacgtc gccgtcaatg actccgggtg gcaccagctc gtcagccact	1620
ggctgaacac tcacgcggtg atggagccgt tcgtgatctc gacgaaccgg caccttagcg	1680
tgacgcaccc ggtgcacaag ctgctgagcc cgcactaccg cgacaccatg accatcaacg	1740
cgtggcgcg gcagacgtc atcaacgccg gcggcatctt cgagatgacg gtgttcccgg	1800
gcaagttcgc gttggggatg tcggccgtgg tgtacaagga ctggaagttc accgagcagg	1860
gactgccgga cgatctcatc aagaggggca tggcggtgga ggaccgctc agcccgtaca	1920
aggtgcggtt gctggtgtcg gactacctgt acgcggcgga cgggctggcg atctggcacg	1980
ccattgagca gtacgtgagc gactacctgt ccatctacta cccgaacgac ggcgtgctgc	2040
agggcgatac ggaggtgcag gcgtggtgga aggagacgcg cgaggtcggg cacggcgacc	2100
tcaaggacgc cccatggtgg cccaagatgc aaagtgtgcc ggagctggcc aaggcgtgca	2160
ccaccatcat ctggatcggg tcggcgtgc atgcggcagt caacttcggg cagtaccctt	2220
acgcggggtt cctcccgaac cggccgacgg tgagccggcg ccgcatgccg gagcccggca	2280
cggaggagta cgcggagctg gagcgcgacc cggagcgggc cttcatccac accatcacga	2340
gccagatcca gaccatcatc ggcgtgtcgc tgctggaggt gctgtcgaag cactcctccg	2400

## SEQLIST.ST25

```

acgagctgta cctcgggag cgggacacgc cggagtggac ctcggaccca aaggccctgg 2460
aggtgttcaa gcggttcagc gaccggctgg tggagatcga gagcaagggtg gtgggcatga 2520
accatgaccc ggagctcaag aaccgcaacg gcccggctaa gtttcctac atgctgctct 2580
acccaacac ctccgaccac aagggcgccg ctgccgggct taccgccaag ggcattccca 2640
acagcatctc catctaattc aagccatcgg caaccatgga tgaataaagg gcgttcgcca 2700
cgtacgaaac ttgtcgagag attggtgtag tgtgtgtctg tgacagtact atgtcagcag 2760
ttgctcttta agccgaataa ataaagcaga tttgcttcca aaaaaaaaaa aaaaaaaa 2818

```

&lt;210&gt; 11

&lt;211&gt; 4663

&lt;212&gt; DNA

&lt;213&gt; Hordeum vulgare

&lt;220&gt;

&lt;221&gt; variation

&lt;222&gt; (2346)..(2348)

&lt;223&gt; "n" is a, c, t, or g encoding an acidic, basic, or polar amino acid

&lt;400&gt; 11

```

cagcccatg catgcacatg cacatgcaca tgcacatgca gtgcagccaa gcaccgctcg 60
atgggcatc acccgtcacg ggaccggagc gcgccatgcg aagcacgagg agggcacgtc 120
accgtccgag cgcagcacgt ggagagcacg tcgccgtccg atccatctct ccaaagccga 180
gcgccacacc accgggaccg gaccgggacc ggcctataaa ttgcccggac cgagctgcaa 240
gcagctcctc acacacactc acgcaacaca catccatctt cactgaaaag tgaaaaacag 300
tgtgtggtg ccattggtg gagcagtga aagcaggaga ggaggccaag aacaagatgc 360
tgctgggagg gctgatcgac accctcacgg gggcgaacaa gagcgcccgg ctcaagggca 420
cgggtggtgct catgcgcaag aacgtgctgg acctcaacga cttcggcgcc accatcatcg 480
acggcatcgg cgagttcctc ggcaagggtg tcacctgcca gcttatcagc tccaccgccc 540
tcgaccaagg taatcactac cctcctccgg ccttctcctc tgtttacaag atatagtatt 600
tctttcgtgt gggccggcgg ccatggatgg atggatgtgt ctggatcggc taaagaagat 660
aggatagcta gccctggccg gtcgtcttta cctgagcatg ggcataatgcc atcgaaaaaa 720
gagacaacag catgcatgca tgggtgcgagc accagaccac gcagagcacc ggatgctcga 780

```

## ..SEQLIST.ST25

gacaaagcaa cacaacaagc aaggacgaca cgtcaaaaagc aacacaacaa gcaaggacgg	840
cacgtcaaaa gcaacacaaa cctaaactaa agcacaaaga cgtaagagca agcacacaat	900
cagcaggcta taaacagttg tcatcaaaaa caacgctgga agagagagag aaggaaggaa	960
gtagtagcca tgaaaaatta aatcaccggg cgttgctctt tgcccaacaa ttaatcaagc	1020
agggtacgtg gcatgtatag ttcttgtaag taaactaagc atgtgatatg agaaggtagc	1080
tggtggtgca gacaacggcg gtcgcgggaa ggtgggcgcg gaggcggagc tggagcagtg	1140
ggtgacgagc ctgccgtcgc tgacgacggg ggagtccaag ttcggcctca ccttcgactg	1200
ggaggtggag aagctcgggg tgccgggcgc catcgtcgtc aacaactacc acagctccga	1260
gttcctgctt aaaaccatca cctccacga cgtccccggc cgcagcggca acctcacctt	1320
cgtcgccaac tcatggatct accccgccgc caactaccga tacagccgcg tcttcttcgc	1380
caacgacgtg cgtggatttt cctctacttt cctctccttt cattttcacc gccttcgtca	1440
ttcatggtcg atcattaagt cttgccagga caatagatga tgagctagga gtggttacca	1500
cttagcagta cgtacattat ttattccgtg ttggtagaaa aggatatggt ttggtgcaga	1560
tcgacacaag attgaatgaa agttgcaccg tggcaccgtg gcagcgtggt aggtgaaaat	1620
aactgttgca cggatccacc cacatgattg ttttcatgaa taaacttttt aaggatgtgt	1680
ctagccacat ctagatgcat gtcacataat tattgcatac caaaacgatt aaattaagca	1740
taaaaagaaa aggaaaaaaa tactcacata tctcgacgta agatcaatga tatagtattt	1800
agatatgcaa tatttatctt acatctaaac ctttcttcat tcttaaataat aagacatttg	1860
taagatttca ctatggacaa catacgaaac aaaatcagtg gatctctcta tgcattcatt	1920
atgtagtcta taataaaatc tttaaaagat cgtatatattt gcaacggagg gagtaaaaca	1980
taacttttta atagtaatgt tgcacggctc cacactcgca gacgtacctg ccgagccaga	2040
tgccggcggc gctgaagccg taccgcgacg acgagctccg gaacctgcgt ggcgacgacc	2100
agcagggccc gtaccaggag cagcaccgca tctaccgcta cgacgtctac aacgacctcg	2160
gcgagggccc ccccatcctc ggcggaact ccgaccacc ttacctcgc cgcgccgca	2220
cggagcgcaa gcccacgcc agcgaccgca gcctggagag cgggtgtcg ctgctggagc	2280
agatctacgt gccgcgggac gagaagttcg gccacctcaa gacgtccgac ttcctgggct	2340
actccatcaa ggccatcacg cagggcatcc tgccggccgt gcgcacctac gtggacacca	2400
ccccggcg gttcgactcc ttccaggaca tcatcaacct ctatgagggc ggcatcaagc	2460
tgccaaaggt ggccgcctcg gaggagctcc gtaagcagtt cccgctccag ctcatcaagg	2520
acctcctccc cgtcggcggc gactccctgc ttaagctccc cgtgccccac atcatccagg	2580
agaacaagca ggcgtggagg accgacgagg agttcgcacg ggaggtgctc gccnnngtca	2640
acccggtcat gatcacgcgt ctcacggtga gtcagcgatt atttgttcat tgtgtgtgta	2700



## SEQLIST.ST25

tggtgtccat ggtgagaaag tgcagatctt gatttgcggt gggtcgcatg cacgcatgct	2760
gcatgcatgc aggagttccc gccaaaaagt agtctggacc ctagcaagtt tggtgaccac	2820
accagcacca tcacggcgga gcacatagag aagaacctcg agggcctcac ggtgcagcag	2880
gtaattggtc caagccatcg acatcaacta tgatttacct aggagtaatt ggtagctgta	2940
gataatttgg ctctgttgca attaatgtga tgctggccga tcaagtgatc gtattgggtt	3000
tgaaatttgc aggcgctgga aagcaacagg ctgtacatcc ttgatcacca tgaccggttc	3060
atgccgttcc tgatcgacgt caacaacctg cccggcaact tcattctacgc cacgaggacc	3120
ctcttcttcc tgcgcggcga cggcaggctc acgccgctcg ccatcgagct gagcgagccc	3180
atcatccagg gcggccttac cacggccaag agcaaggttt acacgccggt gccagcggc	3240
tccgtcgaag gctgggtgtg ggagctcgcc aaggcctacg tcgccgtcaa tgactccggg	3300
tggcaccagc tcgtcagcca ctggtacgtt ctccacggtc gatgtgattc agtcagtcga	3360
tgcacaacaa ctgatcgaaa tatgattgat tgaaacgcgc aggtgaaca ctcacgcggt	3420
gatggagccg ttcgtgatct cgacgaaccg gcaccttagc gtgacgcacc cgggtgcacaa	3480
gctgctgagc ccgcactacc gcgacaccat gaccatcaac gcgctggcgc ggcagacgct	3540
catcaacgcc ggcggcatct tcgagatgac ggtgttcccg ggcaagtctg cgttggggat	3600
gtcggccgtg gtgtacaagg actggaagtt caccgagcag ggactgccgg acgatctcat	3660
caagaggtag gtacctggtg aatgttatga atgtgtaaaa caaattgggc gtctcgctca	3720
ctgacaggaa cgttggtaaaa aaaatgcagg ggcattggcg tggaggacct gtcgagcccc	3780
tacaaggtag ggttgctggt gtcggactac ccgtacgcg cgacggggt ggcgatctgg	3840
cacgccattg agcagtacgt gagcgagtac ctggccatct actaccgaa cgacggcgtg	3900
ctgcaggcgc atacggaggt gcaggcgtg tggaaggaga gcgcgaggt cgggcacggc	3960
gacctcaagg acgccccatg gtggcccaag atgcaaagtg tgccggagct ggccaaggcg	4020
tgcaccacca tcattctggat cgggtcggcg ctgcatgcgg cagtcaactt cgggcagtac	4080
ccctacgcgg ggttctctcc gaaccggccg acggtgagcc ggcgccgat gccggagccc	4140
ggcacggagg agtacgcgga gctggagcgc gacccggagc gggccttcat ccacaccatc	4200
acgagccaga tccagaccat catcggcgtg tcgctgctgg aggtgctgtc gaagcactcc	4260
tccgacgagc tgtacctcgg gcagcgggac acgccggagt ggacctcga cccaaaggcc	4320
ctggagggtg tcaagcggtt cagcgaccgg ctggtggaga tcgagagcaa ggtggtgggc	4380
atgaaccatg acccggagct caagaaccgc aacggcccg ctaagtttcc ctacatgctg	4440
ctctacccca acacctccga ccacaagggc gccgctgccg ggcttaccgc caagggcatc	4500
cccaacagca tctccatcta atctaagcca tcggcaacca tggatgaata aagggcgttc	4560

## SEQLIST.ST25

gccacgtacg aaacttgctg agagattggt gtagtgtgtg tctgtgacag tactatgtca 4620  
gcagttgctc ttttaagccga ataaataaag cagatttgct tcc 4663

<210> 12

<211> 862

<212> PRT

<213> Hordeum vulgare

<220>

<221> variation

<222> (368)..(368)

<223> "Xaa" is an acidic, basic, or polar amino acid

<400> 12

Met Leu Leu Gly Gly Leu Ile Asp Thr Leu Thr Gly Ala Asn Lys Ser  
1 5 10 15  
Ala Arg Leu Lys Gly Thr Val Val Leu Met Arg Lys Asn Val Leu Asp  
20 25 30  
Leu Asn Asp Phe Gly Ala Thr Ile Ile Asp Gly Ile Gly Glu Phe Leu  
35 40 45  
Gly Lys Gly Val Thr Cys Gln Leu Ile Ser Ser Thr Ala Val Asp Gln  
50 55 60  
Asp Asn Gly Gly Arg Gly Lys Val Gly Ala Glu Ala Glu Leu Glu Gln  
65 70 75 80  
Trp Val Thr Ser Leu Pro Ser Leu Thr Thr Gly Glu Ser Lys Phe Gly  
85 90 95  
Leu Thr Phe Asp Trp Glu Val Glu Lys Leu Gly Val Pro Gly Ala Ile  
100 105 110  
Val Val Asn Asn Tyr His Ser Ser Glu Phe Leu Leu Lys Thr Ile Thr  
115 120 125  
Leu His Asp Val Pro Gly Arg Ser Gly Asn Leu Thr Phe Val Ala Asn  
130 135 140  
Ser Trp Ile Tyr Pro Ala Ala Asn Tyr Arg Tyr Ser Arg Val Phe Phe  
145 150 155 160  
Ala Asn Asp Thr Tyr Leu Pro Ser Gln Met Pro Ala Ala Leu Lys Pro  
165 170 175  
Tyr Arg Asp Asp Glu Leu Arg Asn Leu Arg Gly Asp Asp Gln Gln Gly  
180 185 190

## SEQLIST.ST25

Pro Tyr Gln Glu His Asp Arg Ile Tyr Arg Tyr Asp Val Tyr Asn Asp  
 195 200 205  
 Leu Gly Glu Gly Arg Pro Ile Leu Gly Gly Asn Ser Asp His Pro Tyr  
 210 215 220  
 Pro Arg Arg Gly Arg Thr Glu Arg Lys Pro Asn Ala Ser Asp Pro Ser  
 225 230 235 240  
 Leu Glu Ser Arg Leu Ser Leu Leu Glu Gln Ile Tyr Val Pro Arg Asp  
 245 250 255  
 Glu Lys Phe Gly His Leu Lys Thr Ser Asp Phe Leu Gly Tyr Ser Ile  
 260 265 270  
 Lys Ala Ile Thr Gln Gly Ile Leu Pro Ala Val Arg Thr Tyr Val Asp  
 275 280 285  
 Thr Thr Pro Gly Glu Phe Asp Ser Phe Gln Asp Ile Ile Asn Leu Tyr  
 290 295 300  
 Glu Gly Gly Ile Lys Leu Pro Lys Val Ala Ala Leu Glu Glu Leu Arg  
 305 310 315 320  
 Lys Gln Phe Pro Leu Gln Leu Ile Lys Asp Leu Leu Pro Val Gly Gly  
 325 330 335  
 Asp Ser Leu Leu Lys Leu Pro Val Pro His Ile Ile Gln Glu Asn Lys  
 340 345 350  
 Gln Ala Trp Arg Thr Asp Glu Glu Phe Ala Arg Glu Val Leu Ala Xaa  
 355 360 365  
 Val Asn Pro Val Met Ile Thr Arg Leu Thr Glu Phe Pro Pro Lys Ser  
 370 375 380  
 Ser Leu Asp Pro Ser Lys Phe Gly Asp His Thr Ser Thr Ile Thr Ala  
 385 390 395 400  
 Glu His Ile Glu Lys Asn Leu Glu Gly Leu Thr Val Gln Gln Ala Leu  
 405 410 415  
 Glu Ser Asn Arg Leu Tyr Ile Leu Asp His His Asp Arg Phe Met Pro  
 420 425 430  
 Phe Leu Ile Asp Val Asn Asn Leu Pro Gly Asn Phe Ile Tyr Ala Thr  
 435 440 445  
 Arg Thr Leu Phe Phe Leu Arg Gly Asp Gly Arg Leu Thr Pro Leu Ala  
 450 455 460  
 Ile Glu Leu Ser Glu Pro Ile Ile Gln Gly Gly Leu Thr Thr Ala Lys  
 465 470 475 480  
 Ser Lys Val Tyr Thr Pro Val Pro Ser Gly Ser Val Glu Gly Trp Val  
 485 490 495  
 Trp Glu Leu Ala Lys Ala Tyr Val Ala Val Asn Asp Ser Gly Trp His  
 500 505 510  
 Gln Leu Val Ser His Trp Leu Asn Thr His Ala Val Met Glu Pro Phe  
 515 520 525

## SEQLIST.ST25

Val Ile Ser Thr Asn Arg His Leu Ser Val Thr His Pro Val His Lys  
 530 535 540  
 Leu Leu Ser Pro His Tyr Arg Asp Thr Met Thr Ile Asn Ala Leu Ala  
 545 550 555 560  
 Arg Gln Thr Leu Ile Asn Ala Gly Gly Ile Phe Glu Met Thr Val Phe  
 565 570 575  
 Pro Gly Lys Phe Ala Leu Gly Met Ser Ala Val Val Tyr Lys Asp Trp  
 580 585 590  
 Lys Phe Thr Glu Gln Gly Leu Pro Asp Asp Leu Ile Lys Arg Gly Met  
 595 600 605  
 Ala Val Glu Asp Pro Ser Ser Pro Tyr Lys Val Arg Leu Leu Val Ser  
 610 615 620  
 Asp Tyr Pro Tyr Ala Ala Asp Gly Leu Ala Ile Trp His Ala Ile Glu  
 625 630 635 640  
 Gln Tyr Val Ser Glu Tyr Leu Ala Ile Tyr Tyr Pro Asn Asp Gly Val  
 645 650 655  
 Leu Gln Gly Asp Thr Glu Val Gln Ala Trp Trp Lys Glu Thr Arg Glu  
 660 665 670  
 Val Gly His Gly Asp Leu Lys Asp Ala Pro Trp Trp Pro Lys Met Gln  
 675 680 685  
 Ser Val Pro Glu Leu Ala Lys Ala Cys Thr Thr Ile Ile Trp Ile Gly  
 690 695 700  
 Ser Ala Leu His Ala Ala Val Asn Phe Gly Gln Tyr Pro Tyr Ala Gly  
 705 710 715 720  
 Phe Leu Pro Asn Arg Pro Thr Val Ser Arg Arg Arg Met Pro Glu Pro  
 725 730 735  
 Gly Thr Glu Glu Tyr Ala Glu Leu Glu Arg Asp Pro Glu Arg Ala Phe  
 740 745 750  
 Ile His Thr Ile Thr Ser Gln Ile Gln Thr Ile Ile Gly Val Ser Leu  
 755 760 765  
 Leu Glu Val Leu Ser Lys His Ser Ser Asp Glu Leu Tyr Leu Gly Gln  
 770 775 780  
 Arg Asp Thr Pro Glu Trp Thr Ser Asp Pro Lys Ala Leu Glu Val Phe  
 785 790 795 800  
 Lys Arg Phe Ser Asp Arg Leu Val Glu Ile Glu Ser Lys Val Val Gly  
 805 810 815  
 Met Asn His Asp Pro Glu Leu Lys Asn Arg Asn Gly Pro Ala Lys Phe  
 820 825 830  
 Pro Tyr Met Leu Leu Tyr Pro Asn Thr Ser Asp His Lys Gly Ala Ala  
 835 840 845  
 Ala Gly Leu Thr Ala Lys Gly Ile Pro Asn Ser Ile Ser Ile  
 850 855 860

## SEQLIST.ST25

&lt;210&gt; 13

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 13

cgctacgacg tctacaacga

20

&lt;210&gt; 14

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 14

cagactactt tttggcggga

20

&lt;210&gt; 15

&lt;211&gt; 839

&lt;212&gt; PRT

&lt;213&gt; Glycine max

&lt;400&gt; 15

Met Phe Ser Ala Gly His Lys Ile Lys Gly Thr Val Val Leu Met Pro  
1 5 10 15Lys Asn Glu Leu Glu Val Asn Pro Asp Gly Ser Ala Val Asp Asn Leu  
20 25 30Asn Ala Phe Leu Gly Arg Ser Val Ser Leu Gln Leu Ile Ser Ala Thr  
35 40 45Lys Ala Asp Ala His Gly Lys Gly Lys Val Gly Lys Asp Thr Phe Leu  
50 55 60Glu Gly Ile Asn Thr Ser Leu Pro Thr Leu Gly Ala Gly Glu Ser Ala  
65 70 75 80Phe Asn Ile His Phe Glu Trp Asp Gly Ser Met Gly Ile Pro Gly Ala  
85 90 95Phe Tyr Ile Lys Asn Tyr Met Gln Val Glu Phe Phe Leu Lys Ser Leu  
100 105 110Thr Leu Glu Ala Ile Ser Asn Gln Gly Thr Ile Arg Phe Val Cys Asn  
115 120 125

## SEQLIST.ST25

Ser Trp Val Tyr Asn Thr Lys Leu Tyr Lys Ser Val Arg Ile Phe Phe  
 130 135 140  
 Ala Asn His Thr Tyr Val Pro Ser Glu Thr Pro Ala Pro Leu Val Ser  
 145 150 155 160  
 Tyr Arg Glu Glu Glu Leu Lys Ser Leu Arg Gly Asn Gly Thr Gly Glu  
 165 170 175  
 Arg Lys Glu Tyr Asp Arg Ile Tyr Asp Tyr Asp Val Tyr Asn Asp Leu  
 180 185 190  
 Gly Asn Pro Asp Lys Ser Glu Lys Leu Ala Arg Pro Val Leu Gly Gly  
 195 200 205  
 Ser Ser Thr Phe Pro Tyr Pro Arg Arg Gly Arg Thr Gly Arg Gly Pro  
 210 215 220  
 Thr Val Thr Asp Pro Asn Thr Glu Lys Gln Gly Glu Val Phe Tyr Val  
 225 230 235 240  
 Pro Arg Asp Glu Asn Leu Gly His Leu Lys Ser Lys Asp Ala Leu Glu  
 245 250 255  
 Ile Gly Thr Lys Ser Leu Ser Gln Ile Val Gln Pro Ala Phe Glu Ser  
 260 265 270  
 Ala Phe Asp Leu Lys Ser Thr Pro Ile Glu Phe His Ser Phe Gln Asp  
 275 280 285  
 Val His Asp Leu Tyr Glu Gly Gly Ile Lys Leu Pro Arg Asp Val Ile  
 290 295 300  
 Ser Thr Ile Ile Pro Leu Pro Val Ile Lys Glu Leu Tyr Arg Thr Asp  
 305 310 315 320  
 Gly Gln His Ile Leu Lys Phe Pro Gln Pro His Val Val Gln Val Ser  
 325 330 335  
 Gln Ser Ala Trp Met Thr Asp Glu Glu Phe Ala Arg Glu Met Ile Ala  
 340 345 350  
 Gly Val Asn Pro Cys Val Ile Arg Gly Leu Glu Glu Phe Pro Pro Lys  
 355 360 365  
 Ser Asn Leu Asp Pro Ala Ile Tyr Gly Asp Gln Ser Ser Lys Ile Thr  
 370 375 380  
 Ala Asp Ser Leu Asp Leu Asp Gly Tyr Thr Met Asp Glu Ala Leu Gly  
 385 390 395 400  
 Ser Arg Arg Leu Phe Met Leu Asp Tyr His Asp Ile Phe Met Pro Tyr  
 405 410 415  
 Val Arg Gln Ile Asn Gln Leu Asn Ser Ala Lys Thr Tyr Ala Thr Arg  
 420 425 430  
 Thr Ile Leu Phe Leu Arg Glu Asp Gly Thr Leu Lys Pro Val Ala Ile  
 435 440 445  
 Glu Leu Ser Leu Pro His Ser Ala Gly Asp Leu Ser Ala Ala Val Ser  
 450 455 460

## SEQLIST.ST25

Gln Val Val Leu Pro Ala Lys Glu Gly Val Glu Ser Thr Ile Trp Leu  
 465 470 475 480  
 Leu Ala Lys Ala Tyr Val Ile Val Asn Asp Ser Cys Tyr His Gln Leu  
 485 490 495  
 Met Ser His Trp Leu Asn Thr His Ala Ala Met Glu Pro Phe Val Ile  
 500 505 510  
 Ala Thr His Arg His Leu Ser Val Leu His Pro Ile Tyr Lys Leu Leu  
 515 520 525  
 Thr Pro His Tyr Arg Asn Asn Met Asn Ile Asn Ala Leu Ala Arg Gln  
 530 535 540  
 Ser Leu Ile Asn Ala Asn Gly Ile Ile Glu Thr Thr Phe Leu Pro Ser  
 545 550 555 560  
 Lys Tyr Ser Val Glu Met Ser Ser Ala Val Tyr Lys Asn Trp Val Phe  
 565 570 575  
 Thr Asp Gln Ala Leu Pro Ala Asp Leu Ile Lys Arg Gly Val Ala Ile  
 580 585 590  
 Lys Asp Pro Ser Thr Pro His Gly Val Arg Leu Leu Ile Glu Asp Tyr  
 595 600 605  
 Pro Tyr Ala Ala Asp Gly Leu Glu Ile Trp Ala Ala Ile Lys Thr Trp  
 610 615 620  
 Val Gln Glu Tyr Val Pro Leu Tyr Tyr Ala Arg Asp Asp Asp Val Lys  
 625 630 635 640  
 Asn Asp Ser Glu Leu Gln His Trp Trp Lys Glu Ala Val Glu Lys Gly  
 645 650 655  
 His Gly Asp Leu Lys Asp Lys Pro Trp Trp Pro Lys Leu Gln Thr Leu  
 660 665 670  
 Glu Asp Leu Val Glu Val Cys Leu Ile Ile Ile Trp Ile Ala Ser Ala  
 675 680 685  
 Leu His Ala Ala Val Asn Phe Gly Gln Tyr Pro Tyr Gly Gly Leu Ile  
 690 695 700  
 Met Asn Arg Pro Thr Ala Ser Arg Arg Leu Leu Pro Glu Lys Gly Thr  
 705 710 715 720  
 Pro Glu Tyr Glu Glu Met Ile Asn Asn His Glu Lys Ala Tyr Leu Arg  
 725 730 735  
 Thr Ile Thr Ser Lys Leu Pro Thr Leu Ile Ser Leu Ser Val Ile Glu  
 740 745 750  
 Ile Leu Ser Thr His Ala Ser Asp Glu Val Tyr Leu Gly Gln Arg Asp  
 755 760 765  
 Asn Pro His Trp Thr Ser Asp Ser Lys Ala Leu Gln Ala Phe Gln Lys  
 770 775 780  
 Phe Gly Asn Lys Leu Lys Glu Ile Glu Glu Lys Leu Val Arg Arg Asn  
 785 790 795 800

## SEQLIST.ST25

Asn Asp Pro Ser Leu Gln Gly Asn Arg Leu Gly Pro Val Gln Leu Pro  
 805 810 815

Tyr Thr Leu Leu Tyr Pro Ser Ser Glu Gly Leu Thr Phe Arg Gly  
 820 825 830

Ile Pro Asn Ser Ile Ser Ile  
 835

<210> 16

<211> 865

<212> PRT

<213> Glycine max

<400> 16

Met Phe Ser Val Pro Gly Val Ser Gly Ile Leu Asn Arg Gly Gly Gly  
 1 5 10 15

His Lys Ile Lys Gly Thr Val Val Leu Met Arg Lys Asn Val Leu Asp  
 20 25 30

Phe Asn Ser Val Ala Asp Leu Thr Lys Gly Asn Val Gly Gly Leu Ile  
 35 40 45

Gly Thr Gly Leu Asn Val Val Gly Ser Thr Leu Asp Asn Leu Thr Ala  
 50 55 60

Phe Leu Gly Arg Ser Val Ala Leu Gln Leu Ile Ser Ala Thr Lys Pro  
 65 70 75 80

Leu Ala Asn Gly Lys Gly Lys Val Gly Lys Asp Thr Phe Leu Glu Gly  
 85 90 95

Ile Ile Val Ser Leu Pro Thr Leu Gly Ala Gly Glu Ser Ala Phe Asn  
 100 105 110

Ile Gln Phe Glu Trp Asp Glu Ser Met Gly Ile Pro Gly Ala Phe Tyr  
 115 120 125

Ile Lys Asn Tyr Met Gln Val Glu Phe Tyr Leu Lys Ser Leu Thr Leu  
 130 135 140

Glu Asp Val Pro Asn Gln Gly Thr Ile Arg Phe Val Cys Asn Ser Trp  
 145 150 155 160

Val Tyr Asn Thr Lys Leu Tyr Lys Ser Val Arg Ile Phe Phe Ala Asn  
 165 170 175

His Thr Tyr Val Pro Ser Glu Thr Pro Ala Ala Leu Val Gly Tyr Arg  
 180 185 190

Glu Glu Glu Leu Lys Asn Leu Arg Gly Asp Gly Lys Gly Glu Arg Lys  
 195 200 205

Glu His Asp Arg Ile Tyr Asp Tyr Asp Val Tyr Asn Asp Leu Gly Asn  
 210 215 220



## SEQLIST.ST25

Pro Asp His Gly Glu Asn Phe Ala Arg Pro Ile Leu Gly Gly Ser Ser  
 225 230 235 240  
 Thr His Pro Tyr Pro Arg Arg Gly Arg Thr Gly Arg Tyr Pro Thr Arg  
 245 250 255  
 Lys Asp Gln Asn Ser Glu Lys Pro Gly Glu Val Tyr Val Pro Arg Asp  
 260 265 270  
 Glu Asn Phe Gly His Leu Lys Ser Ser Asp Phe Leu Ala Tyr Gly Ile  
 275 280 285  
 Lys Ser Leu Ser Gln Tyr Val Leu Pro Ala Phe Glu Ser Val Phe Asp  
 290 295 300  
 Leu Asn Phe Thr Pro Asn Glu Phe Asp Ser Phe Gln Asp Val Arg Asp  
 305 310 315 320  
 Leu His Glu Gly Gly Ile Lys Leu Pro Thr Glu Val Ile Ser Thr Ile  
 325 330 335  
 Met Pro Leu Pro Val Val Lys Glu Leu Phe Arg Thr Asp Gly Glu Gln  
 340 345 350  
 Val Leu Lys Phe Pro Pro Pro His Val Ile Gln Val Ser Lys Ser Ala  
 355 360 365  
 Trp Met Thr Asp Glu Glu Phe Ala Arg Glu Met Val Ala Gly Val Asn  
 370 375 380  
 Pro Cys Val Ile Arg Gly Leu Gln Glu Phe Pro Pro Lys Ser Asn Leu  
 385 390 395 400  
 Asp Pro Thr Ile Tyr Gly Glu Gln Thr Ser Lys Ile Thr Ala Asp Ala  
 405 410 415  
 Leu Asp Leu Asp Gly Tyr Thr Val Asp Glu Ala Leu Ala Ser Arg Arg  
 420 425 430  
 Leu Phe Met Leu Asp Tyr His Asp Val Phe Met Pro Tyr Ile Arg Arg  
 435 440 445  
 Ile Asn Gln Thr Tyr Ala Lys Ala Tyr Ala Thr Arg Thr Ile Leu Phe  
 450 455 460  
 Leu Arg Glu Asn Gly Thr Leu Lys Pro Val Ala Ile Glu Leu Ser Leu  
 465 470 475 480  
 Pro His Pro Ala Gly Asp Leu Ser Gly Ala Val Ser Gln Val Ile Leu  
 485 490 495  
 Pro Ala Lys Glu Gly Val Glu Ser Thr Ile Trp Leu Leu Ala Lys Ala  
 500 505 510  
 Tyr Val Val Val Asn Asp Ser Cys Tyr His Gln Leu Met Ser His Trp  
 515 520 525  
 Leu Asn Thr His Ala Val Ile Glu Pro Phe Ile Ile Ala Thr Asn Arg  
 530 535 540  
 His Leu Ser Ala Leu His Pro Ile Tyr Lys Leu Leu Thr Pro His Tyr  
 545 550 555 560

## SEQLIST.ST25

Arg Asp Thr Met Asn Ile Asn Ala Leu Ala Arg Gln Ser Leu Ile Asn  
 565 570 575  
 Ala Asp Gly Ile Ile Glu Lys Ser Phe Leu Pro Ser Lys His Ser Val  
 580 585 590  
 Glu Met Ser Ser Ala Val Tyr Lys Asn Trp Val Phe Thr Asp Gln Ala  
 595 600 605  
 Leu Pro Ala Asp Leu Ile Lys Arg Gly Val Ala Ile Lys Asp Pro Ser  
 610 615 620  
 Ala Pro His Gly Leu Arg Leu Leu Ile Glu Asp Tyr Pro Tyr Ala Val  
 625 630 635 640  
 Asp Gly Leu Glu Ile Trp Ala Ala Ile Lys Thr Trp Val Gln Glu Tyr  
 645 650 655  
 Val Ser Leu Tyr Tyr Ala Arg Asp Asp Asp Val Lys Pro Asp Ser Glu  
 660 665 670  
 Leu Gln Gln Trp Trp Lys Glu Ala Val Glu Lys Gly His Gly Asp Leu  
 675 680 685  
 Lys Asp Lys Pro Trp Trp Pro Lys Leu Gln Thr Ile Glu Glu Leu Val  
 690 695 700  
 Glu Ile Cys Thr Ile Ile Ile Trp Thr Ala Ser Ala Leu His Ala Ala  
 705 710 715 720  
 Val Asn Phe Gly Gln Tyr Pro Tyr Gly Gly Phe Ile Leu Asn Arg Pro  
 725 730 735  
 Thr Ser Ser Arg Arg Leu Leu Pro Glu Lys Gly Thr Pro Glu Tyr Glu  
 740 745 750  
 Glu Met Val Lys Ser His Gln Lys Ala Tyr Leu Arg Thr Ile Thr Ser  
 755 760 765  
 Lys Phe Gln Thr Leu Val Asp Leu Ser Val Ile Glu Ile Leu Ser Arg  
 770 775 780  
 His Ala Ser Asp Glu Val Tyr Leu Gly Gln Arg Asp Asn Pro His Trp  
 785 790 795 800  
 Thr Ser Asp Ser Lys Ala Leu Gln Ala Phe Gln Lys Phe Gly Asn Lys  
 805 810 815  
 Leu Lys Glu Ile Glu Glu Lys Leu Ala Arg Lys Asn Asn Asp Gln Ser  
 820 825 830  
 Leu Ser Asn Arg Leu Gly Pro Val Gln Leu Pro Tyr Thr Leu Leu His  
 835 840 845  
 Pro Asn Ser Glu Gly Leu Thr Cys Arg Gly Ile Pro Asn Ser Ile Ser  
 850 855 860  
 Ile  
 865

&lt;210&gt; 17

## SEQLIST.ST25

&lt;211&gt; 857

&lt;212&gt; PRT

&lt;213&gt; Glycine max

&lt;400&gt; 17

Met Leu Gly Gly Leu Leu His Arg Gly His Lys Ile Lys Gly Thr Val  
 1 5 10 15  
 Val Leu Met Arg Lys Asn Val Leu Asp Val Asn Ser Val Thr Ser Val  
 20 25 30  
 Gly Gly Ile Ile Gly Gln Gly Leu Asp Leu Val Gly Ser Thr Leu Asp  
 35 40 45  
 Thr Leu Thr Ala Phe Leu Gly Arg Ser Val Ser Leu Gln Leu Ile Ser  
 50 55 60  
 Ala Thr Lys Ala Asp Ala Asn Gly Lys Gly Lys Leu Gly Lys Ala Thr  
 65 70 75 80  
 Phe Leu Glu Gly Ile Ile Thr Ser Leu Pro Thr Leu Gly Ala Gly Gln  
 85 90 95  
 Ser Ala Phe Lys Ile Asn Phe Glu Trp Asp Asp Gly Ser Gly Ile Pro  
 100 105 110  
 Gly Ala Phe Tyr Ile Lys Asn Phe Met Gln Thr Glu Phe Phe Leu Val  
 115 120 125  
 Ser Leu Thr Leu Glu Asp Ile Pro Asn His Gly Ser Ile His Phe Val  
 130 135 140  
 Cys Asn Ser Trp Ile Tyr Asn Ala Lys Leu Phe Lys Ser Asp Arg Ile  
 145 150 155 160  
 Phe Phe Ala Asn Gln Thr Tyr Leu Pro Ser Glu Thr Pro Ala Pro Leu  
 165 170 175  
 Val Lys Tyr Arg Glu Glu Glu Leu His Asn Leu Arg Gly Asp Gly Thr  
 180 185 190  
 Gly Glu Arg Lys Glu Trp Glu Arg Ile Tyr Asp Tyr Asp Val Tyr Asn  
 195 200 205  
 Asp Leu Gly Asp Pro Asp Lys Gly Glu Asn His Ala Arg Pro Val Leu  
 210 215 220  
 Gly Gly Asn Asp Thr Phe Pro Tyr Pro Arg Arg Gly Arg Thr Gly Arg  
 225 230 235 240  
 Lys Pro Thr Arg Lys Asp Pro Asn Ser Glu Ser Arg Ser Asn Asp Val  
 245 250 255  
 Tyr Leu Pro Arg Asp Glu Ala Phe Gly His Leu Lys Ser Ser Asp Phe  
 260 265 270  
 Leu Thr Tyr Gly Leu Lys Ser Val Ser Gln Asn Val Leu Pro Leu Leu  
 275 280 285

## SEQLIST.ST25

Gln Ser Ala Phe Asp Leu Asn Phe Thr Pro Arg Glu Phe Asp Ser Phe  
 290 295 300  
 Asp Glu Val His Gly Leu Tyr Ser Gly Gly Ile Lys Leu Pro Thr Asp  
 305 310 315 320  
 Ile Ile Ser Lys Ile Ser Pro Leu Pro Val Leu Lys Glu Ile Phe Arg  
 325 330 335  
 Thr Asp Gly Glu Gln Ala Leu Lys Phe Pro Pro Pro Lys Val Ile Gln  
 340 345 350  
 Val Ser Lys Ser Ala Trp Met Thr Asp Glu Glu Phe Ala Arg Glu Met  
 355 360 365  
 Leu Ala Gly Val Asn Pro Asn Leu Ile Arg Cys Leu Lys Asp Phe Pro  
 370 375 380  
 Pro Arg Ser Lys Leu Asp Ser Gln Val Tyr Gly Asp His Thr Ser Gln  
 385 390 395 400  
 Ile Thr Lys Glu His Leu Glu Pro Asn Leu Glu Gly Leu Thr Val Asp  
 405 410 415  
 Glu Ala Ile Gln Asn Lys Arg Leu Phe Leu Leu Asp His His Asp Pro  
 420 425 430  
 Ile Met Pro Tyr Leu Arg Arg Ile Asn Ala Thr Ser Thr Lys Ala Tyr  
 435 440 445  
 Ala Thr Arg Thr Ile Leu Phe Leu Lys Asn Asp Gly Thr Leu Arg Pro  
 450 455 460  
 Leu Ala Ile Glu Leu Ser Leu Pro His Pro Gln Gly Asp Gln Ser Gly  
 465 470 475 480  
 Ala Phe Ser Gln Val Phe Leu Pro Ala Asp Glu Gly Val Glu Ser Ser  
 485 490 495  
 Ile Trp Leu Leu Ala Lys Ala Tyr Val Val Val Asn Asp Ser Cys Tyr  
 500 505 510  
 His Gln Leu Val Ser His Trp Leu Asn Thr His Ala Val Val Glu Pro  
 515 520 525  
 Phe Ile Ile Ala Thr Asn Arg His Leu Ser Val Val His Pro Ile Tyr  
 530 535 540  
 Lys Leu Leu His Pro His Tyr Arg Asp Thr Met Asn Ile Asn Gly Leu  
 545 550 555 560  
 Ala Arg Leu Ser Leu Val Asn Asp Gly Gly Val Ile Glu Gln Thr Phe  
 565 570 575  
 Leu Trp Gly Arg Tyr Ser Val Glu Met Ser Ala Val Val Tyr Lys Asp  
 580 585 590  
 Trp Val Phe Thr Asp Gln Ala Leu Pro Ala Asp Leu Ile Lys Arg Gly  
 595 600 605  
 Met Ala Ile Glu Asp Pro Ser Cys Pro His Gly Ile Arg Leu Val Ile  
 610 615 620

## SEQLIST.ST25

Glu Asp Tyr Pro Tyr Thr Val Asp Gly Leu Glu Ile Trp Asp Ala Ile  
 625 630 635 640  
 Lys Thr Trp Val His Glu Tyr Val Phe Leu Tyr Tyr Lys Ser Asp Asp  
 645 650 655  
 Thr Leu Arg Glu Asp Pro Glu Leu Gln Ala Cys Trp Lys Glu Leu Val  
 660 665 670  
 Glu Val Gly His Gly Asp Lys Lys Asn Glu Pro Trp Trp Pro Lys Met  
 675 680 685  
 Gln Thr Arg Glu Glu Leu Val Glu Ala Cys Ala Ile Ile Ile Trp Thr  
 690 695 700  
 Ala Ser Ala Leu His Ala Ala Val Asn Phe Gly Gln Tyr Pro Tyr Gly  
 705 710 715 720  
 Gly Leu Ile Leu Asn Arg Pro Thr Leu Ser Arg Arg Phe Met Pro Glu  
 725 730 735  
 Lys Gly Ser Ala Glu Tyr Glu Glu Leu Arg Lys Asn Pro Gln Lys Ala  
 740 745 750  
 Tyr Leu Lys Thr Ile Thr Pro Lys Phe Gln Thr Leu Ile Asp Leu Ser  
 755 760 765  
 Val Ile Glu Ile Leu Ser Arg His Ala Ser Asp Glu Val Tyr Leu Gly  
 770 775 780  
 Glu Arg Asp Asn Pro Asn Trp Thr Ser Asp Thr Arg Ala Leu Glu Ala  
 785 790 795 800  
 Phe Lys Arg Phe Gly Asn Lys Leu Ala Gln Ile Glu Asn Lys Leu Ser  
 805 810 815  
 Glu Arg Asn Asn Asp Glu Lys Leu Arg Asn Arg Cys Gly Pro Val Gln  
 820 825 830  
 Met Pro Tyr Thr Leu Leu Leu Pro Ser Ser Lys Glu Gly Leu Thr Phe  
 835 840 845  
 Arg Gly Ile Pro Asn Ser Ile Ser Ile  
 850 855

&lt;210&gt; 18

&lt;211&gt; 864

&lt;212&gt; PRT

&lt;213&gt; Hordeum vulgare

&lt;400&gt; 18

Met Leu Gly Val Gly Gly Ile Val Ser Asp Leu Thr Gly Gly Ile Arg  
 1 5 10 15  
 Gly Ala His Leu Lys Gly Ser Val Val Leu Met Arg Lys Asn Ala Leu  
 20 25 30

## SEQLIST.ST25

Asp Phe Asn Asp Phe Gly Ala His Val Met Asp Gly Val Thr Glu Leu  
 35 40 45  
 Leu Gly Arg Gly Val Thr Cys Gln Leu Ile Ser Ser Thr Asn Val Asp  
 50 55 60  
 His Asn Asn Gly Gly Arg Gly Lys Val Gly Ala Glu Ala Asn Leu Glu  
 65 70 75 80  
 Gln Trp Leu Leu Pro Thr Asn Leu Pro Phe Ile Thr Thr Gly Glu Asn  
 85 90 95  
 Lys Phe Ala Val Thr Phe Asp Trp Ser Val Asp Lys Leu Gly Val Pro  
 100 105 110  
 Gly Ala Ile Ile Val Lys Asn Asn His Ala Ser Glu Phe Phe Leu Lys  
 115 120 125  
 Thr Ile Thr Leu Asp Asn Val Pro Gly Arg Gly Thr Ile Val Phe Val  
 130 135 140  
 Ala Asn Ser Trp Val Tyr Pro Gln Ala Lys Tyr Arg Tyr Asn Arg Val  
 145 150 155 160  
 Phe Phe Ala Asn Asp Thr Tyr Leu Pro His Gln Met Pro Ala Ala Leu  
 165 170 175  
 Lys Pro Tyr Arg Asp Asp Glu Leu Arg Asn Leu Arg Gly Asp Asp Gln  
 180 185 190  
 Gln Gly Pro Tyr Leu Asp His Asp Arg Val Tyr Arg Tyr Asp Val Tyr  
 195 200 205  
 Asn Asp Leu Gly Asp Ser Arg Asp Val Leu Gly Gly Ser Lys Asp Leu  
 210 215 220  
 Pro Tyr Pro Arg Arg Cys Arg Thr Gly Arg Lys Pro Ser Asp Ser Lys  
 225 230 235 240  
 Pro Asp His Glu Ser Arg Leu Leu Leu Leu Val Gln Asn Val Tyr Val  
 245 250 255  
 Leu Arg Asp Glu Leu Phe Gly His Leu Lys Gln Ser Asp Leu Leu Gly  
 260 265 270  
 Tyr Thr Leu Lys Gly Trp Leu Asp Gly Ile Ile Leu Ala Ile Arg Thr  
 275 280 285  
 Tyr Val Asp Leu Ser Pro Gly Glu Phe Asp Ser Phe Ala Asp Ile Leu  
 290 295 300  
 Lys Leu Tyr Glu Gly Gly Ile Lys Leu Pro Asn Ile Pro Ala Leu Glu  
 305 310 315 320  
 Glu Val Arg Lys Arg Phe Pro Leu Gln Leu Val Lys Asp Leu Ile Pro  
 325 330 335  
 Lys Gly Gly Asp Phe Leu Leu Lys Leu Pro Lys Pro Glu Ile Ile Lys  
 340 345 350  
 Val Asp Gln Lys Ala Trp Met Thr Asp Glu Glu Phe Ala Arg Glu Met  
 355 360 365

## SEQLIST.ST25

Leu Ala Gly Val Asn Pro Met Met Ile Lys Arg Leu Thr Glu Phe Pro  
 370 375 380  
 Pro Lys Ser Thr Leu Asp Pro Ser Lys Tyr Gly Asp His Thr Ser Thr  
 385 390 395 400  
 Met Thr Glu Glu His Val Ala Lys Ser Leu Glu Gly Leu Thr Val Gln  
 405 410 415  
 Gln Ala Leu Ala Gly Asn Arg Leu Tyr Ile Val Asp Gln His Asp Asn  
 420 425 430  
 Leu Met Pro Phe Leu Ile Asp Ile Asn Asn Leu Asp Ala Ser Phe Val  
 435 440 445  
 Tyr Ala Thr Arg Thr Leu Leu Phe Leu Arg Gly Asp Gly Thr Leu Ala  
 450 455 460  
 Pro Val Ala Ile Glu Leu Ser Ser Pro Leu Ile Gln Gly Glu Leu Thr  
 465 470 475 480  
 Thr Ala Lys Ser Ala Val Tyr Thr Pro Gln His Ala Gly Val Glu Gly  
 485 490 495  
 Trp Ile Trp Gln Leu Ala Lys Ala Tyr Ala Ser Val Asn Asp Tyr Gly  
 500 505 510  
 Trp His Gln Leu Ile Ser His Trp Leu Asn Thr His Ala Val Met Glu  
 515 520 525  
 Pro Phe Val Ile Ala Thr Asn Arg Gln Leu Ser Val Thr His Pro Val  
 530 535 540  
 Tyr Lys Leu Leu His Pro His Tyr Arg Asp Thr Met Asn Ile Asn Ala  
 545 550 555 560  
 Arg Ala Arg Gly Leu Leu Ile Asn Ala Gly Gly Val Ile Glu Met Thr  
 565 570 575  
 Val Phe Pro His Lys His Ala Met Pro Met Ser Ser Met Val Tyr Lys  
 580 585 590  
 His Trp Asn Phe Thr Glu Gln Ala Leu Pro Ala Asp Leu Ile Lys Arg  
 595 600 605  
 Gly Met Ala Val Glu Asp Ala Ser Ser Pro His Lys Val Arg Leu Leu  
 610 615 620  
 Ile Lys Asp Tyr Pro Tyr Ala Thr Asp Gly Leu Ala Val Trp Asp Ala  
 625 630 635 640  
 Ile Glu Gln Trp Val Ser Asp Tyr Leu Thr Ile Tyr Tyr Pro Asn Asp  
 645 650 655  
 Gly Val Leu Gln Gly Asp Val Glu Leu Gln Ala Trp Trp Lys Glu Val  
 660 665 670  
 Arg Glu Val Gly His Gly Asp Leu Lys Asp Ala Ala Trp Trp Pro Lys  
 675 680 685  
 Met Gln Thr Val Ala Glu Leu Ile Lys Ala Cys Ala Thr Ile Ile Trp  
 690 695 700

## SEQLIST.ST25

Thr Gly Ser Ala Leu His Ala Ala Val Asn Phe Gly Gln Tyr Pro Tyr  
 705 710 715 720  
 Ser Gly Tyr His Pro Asn Lys Pro Ser Ala Ser Arg Arg Pro Met Pro  
 725 730 735  
 Val Gln Gly Ser Glu Glu Tyr Ala Glu Leu Glu Arg Asp Pro Glu Lys  
 740 745 750  
 Ala Phe Ile Arg Thr Ile Thr Ser Gln Phe His Ala Leu Val Gly Ile  
 755 760 765  
 Ser Leu Met Glu Ile Leu Ser Lys His Ser Ser Asp Glu Val Tyr Leu  
 770 775 780  
 Gly Gln His Asp Thr Pro Ala Trp Thr Ser Asp Ala Lys Ala Leu Glu  
 785 790 795 800  
 Ala Phe Lys Arg Phe Gly Ala Lys Leu Glu Gly Ile Glu Lys Gln Val  
 805 810 815  
 Val Ala Met Asn Ser Asp Pro Gln Leu Lys Asn Arg Thr Gly Pro Ala  
 820 825 830  
 Lys Phe Pro Tyr Met Leu Leu Tyr Pro Asn Thr Ser Asp His Thr Gly  
 835 840 845  
 Gln Ala Glu Gly Leu Thr Ala Arg Gly Ile Pro Asn Ser Ile Ser Ile  
 850 855 860



## INTERNATIONAL SEARCH REPORT

Intern d Application No

PCT/IB 00/02045

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/02 C12N15/82 A01H5/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, FSTA, WPI Data, PAJ, CAB Data, BIOTECHNOLOGY ABS, SCISEARCH  
CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 119, no. 23, 6 December 1993 (1993-12-06) Columbus, Ohio, US; abstract no. 248220m, KOWAKA, M.: "Malting barley improvement for brewing" page 828; column 2; XP002175449 abstract & NIPPON JOZO KYOKAISHI, vol. 88, no. 8, 1993, pages 574-581, --- -/--	1



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*8\* document member of the same patent family

Date of the actual completion of the international search

23 August 2001

Date of mailing of the international search report

06/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Kok, A

## INTERNATIONAL SEARCH REPORT

Inter d Application No

PCT/IB 00/02045

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WAN HENG WANG ET AL: "Molecular basis of a null mutation in soybean lipoxygenase 2: substitution of glutamine for an iron-ligand histidine." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 91, no. 13, 1994, pages 5828-5832, XP002175448 WASHINGTON US abstract	1
X	--- MCELROY D ET AL: "WHAT'S BREWING IN BARLEY BIOTECHNOLOGY?" BIO/TECHNOLOGY, NEW YORK, US, vol. 13, no. 3, 1995, pages 245-249, XP002024712 ISSN: 0733-222X cited in the application page 247, column 2, last paragraph	11
Y	--- DROST B W ET AL: "FLAVOR STABILITY" JOURNAL OF THE AMERICAN SOCIETY OF BREWING CHEMISTS, vol. 48, no. 4, 1990, pages 124-131, XP000926610 ISSN: 0361-0470 cited in the application	11-15
A	the whole document, especially pages 129-130	1,21-27
Y	--- WO 97 13851 A (PURDUE RESEARCH FOUNDATION (US)) 17 April 1997 (1997-04-17) abstract	11-15
A	--- VAN MECHELEN JAN R ET AL: "Molecular characterization of two lipoxygenases from barley." PLANT MOLECULAR BIOLOGY, vol. 39, no. 6, April 1999 (1999-04), pages 1283-1298, XP001015731 ISSN: 0167-4412 abstract; figure 1 -----	1-10

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Present claim 1 relate to a barley plant defined by reference to desirable characteristic or property, namely characterized by a reduction or absence of lipoxygenase activity.

The claims cover all plants having this characteristic or property, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such plants. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the plant by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the subject-matter of claim 8 and claims dependent thereof.

Present claims 6 and 7 relate to an extremely large number of possible barley plants carrying a mutated *lox-1* gene. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only ONE mutated *lox-1* gene.

In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the subject-matter of claim 8 and claims dependent thereof.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

**information on patent family members**

PCT/IB 00/02045

Form PCT/SA/210 (patent family annex) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**